arXiv:0811.4222v1 [math.AP] 26 Nov 2008

Manuscript submitted to
AIMS' Journals
Volume **X**, Number **0X**, XX **200X**

Website: http://AIMsciences.org
pp. **X–XX**

WELL-POSEDNESS FOR ONE-DIMENSIONAL DERIVATIVE NONLINEAR SCHRÖDINGER EQUATIONS

Chengchun Hao

Institute of Mathematics
Academy of Mathematics & Systems Science, CAS
Beijing 100080, P. R. China

(Communicated by Gigliola Staffilani)

Abstract. In this paper, we investigate the one-dimensional derivative nonlinear Schrödinger equations of the form $iu_t - u_{xx} + i\lambda |u|^k u_x = 0$ with non-zero $\lambda \in \mathbb{R}$ and any real number $k \geqslant 5$. We establish the local well-posedness of the Cauchy problem with any initial data in $H^{1/2}$ by using the gauge transformation and the Littlewood-Paley decomposition.

1. **Introduction.** In the present paper, we consider the following Cauchy problem for the derivative nonlinear Schrödinger equation

$$iu_t - u_{xx} + i\lambda |u|^k u_x = 0, \quad (t,x) \in \mathbb{R}^2, \tag{1.1}$$

$$u(0,x) = u_0(x), \tag{1.2}$$

where $u = u(t,x) : \mathbb{R}^2 \to \mathbb{C}$ is a complex-valued wave function, both $\lambda \neq 0$ and $k \geqslant 5$ are real numbers.

A great deal of interesting research has been devoted to the mathematical analysis for the derivative nonlinear Schrödinger equations [3, 4, 6, 7, 8, 9, 10, 11, 13, 18, 21]. In [13], C. E. Kenig, G. Ponce and L. Vega studied the local existence theory for the Cauchy problem of the derivative nonlinear Schrödinger equations

$$iu_t + u_{xx} + f(u, \bar{u}, u_x, \bar{u}_x) = 0, \quad (t,x) \in \mathbb{R}^2,$$

with small data $u(0,x) = u_0(x)$ in $H^{7/2}$ where f is a polynomial having no constant or linear terms with the lowest order term of degree being greater than or equal to 3. Subsequently, it was improved to H^3 by N. Hayashi and T. Ozawa [11].

If the nonlinearity consists mostly of the conjugate wave $\bar{u}$, then it can be done much better. In the case $f = (\bar{u}_x)^k$, A. Grüenrock, in [8], obtained local well-posedness when $s > s_c = 3/2 - 1/(k-1)$, $s \geqslant 1$, and $k \geqslant 2$ was an integer. In particular, the global well-posedness in H^1 is obtained when $f = i(\bar{u}_x)^2$ with the help of the Bourgain spaces (cf. [2, 23]).

2000 *Mathematics Subject Classification.* 35Q55, 35A07.

Key words and phrases. Derivative nonlinear Schrödinger equations, Cauchy problem, local well-posedness.

The author is supported in part by the Scientific Research Startup Special Foundation on Excellent PhD Thesis and Presidential Award of Chinese Academy of Sciences, NSFC (Grant No. 10601061), and the Innovation Funds of AMSS, CAS of China.

In [21], H. Takaoka discussed the derivative nonlinear Schrödinger equation of the form

$$u_t - iu_{xx} + |u|^2 u_x = 0, \quad (t,x) \in \mathbb{R}^2,$$

and obtained the local well-posedness in H^s for $s \geqslant 1/2$ by performing a fixed point argument in an adapted Bourgain space $X_{s,b}$ which yields a C^∞-solution map.

A very similar equation to (1.1) is the generalized Benjamin-Ono equation

$$u_t + \mathcal{H}u_{xx} \pm u^k u_x = 0, \quad (t,x) \in \mathbb{R}^2, \tag{1.3}$$

where u is a real-valued function, $\mathcal{H}$ is the Hilbert transformation defined by

$$\mathcal{H}f(x) = -i \int_{\mathbb{R}} e^{ix\xi} \mathrm{sgn}(\xi) \hat{f}(\xi) d\xi,$$

and $k \geqslant 2$ is an integer, the symbol $\hat{\cdot}$ (or $\mathscr{F}$) denotes the spatial Fourier transform. For this equation, L. Molinet and F. Ribaud [16, 17] obtained the local well-posedness in the Sobolev space H^s for $s > 1/2$ if $k = 2, 4$, $s \geqslant 3/4$ if $k = 3$ and $s \geqslant 1/2$ if $k \geqslant 5$ by using Tao's gauge transformation. In [14], C. E. Kenig and H. Takaoka have shown the global well-posedness for the case $k = 2$ in H^s for $s \geqslant 1/2$ by combining the gauge transformation with a Littlewood-Paley decomposition and following the compactness argument with a priori estimates with the help of the preservation of the Hamiltonian and the L^2-mass.

In the present paper, we shall generalize the above results to the derivative nonlinear Schrödinger equation with $k \geqslant 5$ by using some ideas in [14]. However, we have to reconstruct new and complicated estimates for the case $k \geqslant 5$ which is quite different from the case $k = 2$.

We first state the main result of this paper as follows, though we shall define later the function space X_T at the end of this section.

Theorem 1.1. *For any $u_0 \in H^{1/2}$, there exist a $T = T(\|u_0\|_{H^{1/2}})$ and a unique solution u of* (1.1)-(1.2) *satisfying*

$$u \in C([-T,T]; H^{1/2}) \cap X_T.$$

For convenience, we now introduce some notations. For nonnegative real numbers A, B, we use $A \lesssim B$ to denote $A \leqslant CB$ for some $C > 0$ which is independent of A and B. $A \sim B$ means $A \lesssim B \lesssim A$, and $A \ll B$ denotes $A \leqslant CB$ for some small $C > 0$ which is also independent of A and B.

To give the Littlewood-Paley decomposition, let ψ be a fixed even C^∞ function with a compact support, $\mathrm{supp}\,\psi \subset \{|\xi| < 2\}$, and $\psi(\xi) = 1$ for $|\xi| \leqslant 1$. Define $\varphi(\xi) = \psi(\xi) - \psi(2\xi)$. Let N be a dyadic number of the form $N = 2^j$, $j \in \mathbb{N} \cup \{0\}$ or $N = 0$. Writing $\varphi_N(\xi) = \varphi(\xi/N)$ for $N \geqslant 1$, we define the convolution operator P_N by $P_N u = \check{\varphi}_N * u$, where the symbol $\check{\cdot}$ (or $\mathscr{F}^{-1}$) denotes the spatial Fourier inverse transform. We define the function φ_0 by $\varphi_0(\xi) = 1 - \sum_N \varphi_N(\xi)$ and denote $P_0 u = \check{\varphi}_0 * u$. Then we introduce a spatial Littlewood-Paley decomposition [20]

$$\sum_N P_N = I.$$

Throughout this paper, we often use the Littlewood-Paley theorem (cf. [20, 23])

$$\left\| \left(\sum_N |P_N \phi|^2 \right)^{1/2} \right\|_{L^p} \sim \|\phi\|_{L^p},$$

for $1 < p < \infty$. We also use more general operators $P_{\ll N}$ and $P_{\lesssim N}$ which are defined by

$$P_{\ll N} = \sum_{M \ll N} P_M, \quad P_{\lesssim N} = \sum_{M \lesssim N} P_M,$$

and $P_{\gg N}$, $P_{\gtrsim N}$ and $P_{\sim N}$ which can be defined in a similar way. The Littlewood-Paley operators commute with derivative operators (including $|\nabla|^s$ and $i\partial_t - \partial_{xx}$), the propagator $S(t) = e^{-it\partial_x^2}$, and conjugation operations, are self-adjoint, and are bounded on every Lebesgue space L^p and homogeneous Sobolev space $\dot{H}^s$ if $1 \leqslant p \leqslant \infty$. Furthermore, they obey the following Sobolev and Bernstein estimates for $\mathbb{R}$ with $s \geqslant 0$ and $1 \leqslant p \leqslant \infty$ (which is similar to those of three dimensions [5]):

$$\begin{aligned} \|P_{\geqslant N} f\|_{L^p} &\lesssim N^{-s} \left\| |\nabla|^s P_{\geqslant N} f\right\|_{L^p}, \\ \left\|P_{\leqslant N} |\nabla|^s f\right\|_{L^p} &\lesssim N^{s} \|P_{\leqslant N} f\|_{L^p}, \\ \left\|P_N |\nabla|^{\pm s} f\right\|_{L^p} &\lesssim N^{\pm s} \|P_N f\|_{L^p}, \end{aligned}$$

which can be verified by combining the Bernstein multiplier theorem [1] and the interpolation theorem of Sobolev spaces.

We define the Lebesgue spaces $L_T^q L_x^p$ and $L_x^p L_T^q$ by the norms

$$\|f\|_{L_T^q L_x^p} = \left\| \|f\|_{L_x^p(\mathbb{R})} \right\|_{L_t^q([0,T])}, \quad \|f\|_{L_x^p L_T^q} = \left\| \|f\|_{L_t^q([0,T])} \right\|_{L_x^p(\mathbb{R})}.$$

In particular, we abbreviate $L_T^q L_x^p$ or $L_x^p L_T^q$ as $L_{x,T}^p$ in the case $p = q$.

We also use the elementary inequality [5]

$$\left\| \left(\sum_N |f_N|^2 \right)^{1/2} \right\|_{L_T^q L_x^p} \leqslant \left(\sum_N \|f_N\|_{L_T^q L_x^p}^2 \right)^{1/2},$$

for all $2 \leqslant q, p \leqslant \infty$ and arbitrary functions f_N, and the dual version

$$\left(\sum_N \|f_N\|_{L_T^{q'} L_x^{p'}}^2 \right)^{1/2} \leqslant \left\| \left(\sum_N |f_N|^2 \right)^{1/2} \right\|_{L_T^{q'} L_x^{p'}},$$

where p' is the conjugate number of p given by $1/p + 1/p' = 1$. It is easy to verify that they also hold if we replace the norm $L_T^q L_x^p$ by the norm $L_x^p L_T^q$ in both side of the above inequalities.

Let $\langle \cdot \rangle = (1 + |\cdot|^2)^{1/2}$. We use the fractional differential operators D_x^s and $\langle D_x \rangle^s$ defined by

$$D_x^s f = \mathscr{F}^{-1} |\xi|^s \mathscr{F} f, \quad \langle D_x \rangle^s f = \mathscr{F}^{-1} \langle \xi \rangle^s \mathscr{F} f.$$

Thus, we can introduce the resolution space. For $T > 0$, we define the function space X_T in a similar way as in [14] by

$$X_T := \{u \in \mathscr{S}'((-T,T) \times \mathbb{R}) : \|u\|_{X_T} < \infty\},$$

where

$$\|u\|_{X_T} = \|u\|_{L_T^\infty H^{1/2}} + \left(\sum_N \|\partial_x P_N u\|_{L_x^\infty L_T^2}^2 \right)^{1/2}$$

$$+\left(\sum_N \|P_N u\|^2_{L^2_x L^\infty_T}\right)^{1/2} + \left(\sum_N \left\|\langle D_x\rangle^{\frac{1}{4}} P_N u\right\|^2_{L^4_x L^\infty_T}\right)^{1/2}.$$

2. **Gauge transformation.** We transform the equation (1.1) by introducing the following complex-valued function $v_N : \mathbb{R}^2 \to \mathbb{C}$ for a dyadic number N given by

$$v_N(t,x) = e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u(t,y)|^k dy} P_N u. \tag{2.1}$$

By computation, we have

$$\begin{aligned} i\partial_t v_N - \partial_x^2 v_N =& -i\lambda e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy}\left[P_N(|u|^k u_x) - |P_{\ll N} u|^k P_N u_x\right] \\ &- \frac{i\lambda}{2} e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy} P_N u (i\partial_t - \partial_x^2)\int_{-\infty}^x |P_{\ll N} u(t,y)|^k dy \\ &+ \frac{\lambda^2}{4} e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy} |P_{\ll N} u|^{2k} P_N u. \end{aligned} \tag{2.2}$$

For the second term, we integrate by parts and have

$$\begin{aligned} &(i\partial_t - \partial_x^2)\int_{-\infty}^x |P_{\ll N} u(t,y)|^k dy \\ =& i\int_{-\infty}^x \frac{k}{2}|P_{\ll N} u|^{k-2}\left(\partial_t P_{\ll N} u\overline{P_{\ll N} u} + P_{\ll N} u \partial_t \overline{P_{\ll N} u}\right) dy \\ &- \frac{k}{2}|P_{\ll N} u|^{k-2}\left(\partial_x P_{\ll N} u\overline{P_{\ll N} u} + P_{\ll N} u\partial_x\overline{P_{\ll N} u}\right) \\ =& \int_{-\infty}^x \frac{k}{2}|P_{\ll N} u|^{k-2}\Big(\overline{P_{\ll N} u} P_{\ll N}(u_{xx} - i\lambda |u|^k u_x) \\ &\qquad\qquad -P_{\ll N} u P_{\ll N}(\bar{u}_{xx} + i\lambda|u|^k \bar{u}_x)\Big) dy \\ &- \frac{k}{2}|P_{\ll N} u|^{k-2}\left(P_{\ll N} u_x\overline{P_{\ll N} u} + P_{\ll N} u\overline{P_{\ll N} u_x}\right) \\ =& \int_{-\infty}^x \frac{k(k-2)}{4}|P_{\ll N} u|^{k-4}\left[(\overline{P_{\ll N} u_y} P_{\ll N} u)^2 - (P_{\ll N} u_y \overline{P_{\ll N} u})^2\right] dy \\ &- \int_{-\infty}^x \frac{i\lambda k}{2}|P_{\ll N} u|^{k-2} P_{\ll N}|u|^k (u_x + \bar{u}_x) dy - k|P_{\ll N} u|^{k-2} P_{\ll N} u\overline{P_{\ll N} u_x} \end{aligned}$$

Thus, v_N obeys the following differential-integral equation

$$\begin{aligned} &i\partial_t v_N - \partial_x^2 v_N(t,x) \\ =& -i\lambda e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy}\left[P_N(|u|^k u_x) - |P_{\ll N} u|^k P_N u_x\right] \\ &- \frac{i\lambda k(k-2)}{8} e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy} P_N u \int_{-\infty}^x |P_{\ll N} u|^{k-4} \cdot \\ &\qquad \left[(\overline{P_{\ll N} u_x} P_{\ll N} u)^2 - (P_{\ll N} u_x \overline{P_{\ll N} u})^2\right] dy \\ &- \frac{\lambda^2 k}{4} e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy} P_N u \int_{-\infty}^x |P_{\ll N} u|^{k-2} P_{\ll N}|u|^k (u_x + \bar{u}_x) dy \\ &+ \frac{i\lambda k}{2} e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy} |P_{\ll N} u|^{k-2} P_N u P_{\ll N} u \overline{P_{\ll N} u_x} \\ &+ \frac{\lambda^2}{4} e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy} |P_{\ll N} u|^{2k} P_N u \end{aligned}$$

$$\equiv I_{N,1}(t,x)+I_{N,2}(t,x)+I_{N,3}(t,x)+I_{N,4}(t,x)+I_{N,5}(t,x). \tag{2.3}$$

The equivalent integral equation reads

$$\begin{aligned} v_N(t) =& S(t)e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u_0(y)|^k dy}P_N u_0 \\ & - i\int_0^t S(t-\tau)[I_{N,1}+I_{N,2}+I_{N,3}+I_{N,4}+I_{N,5}](\tau)d\tau. \end{aligned} \tag{2.4}$$

3. **Preliminaries.** In order to prove the a priori estimate for the equation of v_N, we need the linear estimates associated with the one-dimensional Schrödinger equation. We first recall the Strichartz estimates, smoothing effects and maximal function estimates. For the proofs, one can see [13, 14].

Lemma 3.1. *For all $\phi \in \mathscr{S}(\mathbb{R})$, $\theta \in [0,1]$ and $T \in (0,1)$,*

$$\|S(t)\phi\|_{L_T^{\frac{4}{\theta}}L_x^{\frac{2}{1-\theta}}} \lesssim \|\phi\|_{L^2}, \tag{3.1}$$

$$\|S(t)P_N\phi\|_{L_x^{\frac{2}{1-\theta}}L_T^{\frac{2}{\theta}}} \lesssim \langle N\rangle^{\frac{1}{2}-\theta}\|\phi\|_{L^2}, \tag{3.2}$$

$$\|S(t)\phi\|_{L_x^4L_T^\infty} \lesssim \|\phi\|_{\dot{H}^{\frac{1}{4}}}. \tag{3.3}$$

We also need the $L_T^qL_x^p$ and $L_x^pL_T^q$ estimates for the linear operator $f \mapsto \int_0^t S(t-\tau)f(\tau)d\tau$. For the proofs, one can see [14].

Lemma 3.2. *For $f \in \mathscr{S}(\mathbb{R}^2)$, $\theta \in [0,1]$ and $T \in (0,1)$,*

$$\left\|\int_0^t S(t-\tau)f(\tau)d\tau\right\|_{L_T^{\frac{4}{\theta}}L_x^{\frac{2}{1-\theta}}} \lesssim \|f\|_{L_T^{(\frac{4}{\theta})'}L_x^{(\frac{2}{1-\theta})'}}, \tag{3.4}$$

$$\left\|\langle D_x\rangle^{\frac{\theta}{2}}\int_0^t S(t-\tau)f(\tau)d\tau\right\|_{L_T^\infty L_x^2} \lesssim \|f\|_{L_x^{p(\theta)}L_T^{q(\theta)}}, \tag{3.5}$$

$$\left\|\langle D_x\rangle^{\frac{\theta}{2}}\int_0^t S(t-\tau)P_Nf(\tau)d\tau\right\|_{L_x^2L_T^\infty} \lesssim \langle N\rangle^{\frac{1}{2}}\|f\|_{L_x^{p(\theta)}L_T^{q(\theta)}}, \tag{3.6}$$

$$\left\|\langle D_x\rangle^{\frac{\theta}{2}-\frac{1}{4}}\int_0^t S(t-\tau)f(\tau)d\tau\right\|_{L_x^4L_T^\infty} \lesssim \|f\|_{L_x^{p(\theta)}L_T^{q(\theta)}}, \tag{3.7}$$

$$\left\|\langle D_x\rangle^{\frac{1}{2}}\int_0^t S(t-\tau)f(\tau)d\tau\right\|_{L_x^\infty L_T^2} \lesssim \|f\|_{L_T^1L_x^2}, \tag{3.8}$$

$$\left\|\int_0^t S(t-\tau)P_Nf(\tau)d\tau\right\|_{L_x^{\frac{2}{\theta}}L_T^{\frac{2}{1-\theta}}} \lesssim \langle N\rangle^{\frac{1}{2}-\theta}\|f\|_{L_T^1L_x^2}, \tag{3.9}$$

$$\left\|\int_0^t S(t-\tau)f(\tau)d\tau\right\|_{L_x^4L_T^\infty} \lesssim \|f\|_{L_T^1\dot{H}_x^{\frac{1}{4}}}, \tag{3.10}$$

$$\left\|\partial_x\int_0^t S(t-\tau)f(\tau)d\tau\right\|_{L_x^\infty L_T^2} \lesssim \|f\|_{L_x^1L_T^2}, \tag{3.11}$$

where p' is the conjugate number of $p \in [1,\infty]$, i.e. $1/p + 1/p' = 1$, and

$$\frac{1}{p(\theta)} = \frac{3+\theta}{4}, \quad \frac{1}{q(\theta)} = \frac{3-\theta}{4}.$$

Next, we recall the Leibniz' rule for a product of the form $e^{iF}g$ where F is the spatial primitive of some function f. For the proof, we refer to [14, 17].

Lemma 3.3 ([14, Lemma 3.5]). *Let* $\alpha\in(0,1)$, p, p_1, p_2, q, $q_1\in(1,\infty)$, $q_2\in(0,\infty]$ *with* $\frac{1}{p}=\frac{1}{p_1}+\frac{1}{p_2}$, $\frac{1}{q}=\frac{1}{q_1}+\frac{1}{q_2}$, *and let* $F(t,x)=\int_{-\infty}^{x}f(t,y)dy$, *with real-valued function* f. *Then*

$$\left\|D_x^\alpha(e^{iF}g)\right\|_{L_x^pL_T^q}\lesssim\|f\|_{L_x^{p_1}L_T^{q_1}}\|g\|_{L_x^{p_2}L_T^{q_2}}+\|\langle D_x\rangle^\alpha g\|_{L_x^pL_T^q}.$$

4. **Bilinear estimates.**

In this section, we prove the following space-time estimate which is crucial to the proof of the nonlinear estimates.

Proposition 4.1. *Let* $u\in H^\infty$ *and* $p\geqslant 4$ *be a real number. Then we have*

$$\|u\bar{u}_x\|_{L_x^pL_T^2}\lesssim T^{\frac12}\|u\|_{X_T}^2+(1+T^{\frac14}\|u\|_{X_T})\|u\|_{X_T}\|P_{\gg1}u\|_{X_T}. \tag{4.1}$$

Proof. By the Littlewood-Paley decomposition, we can write

$$\begin{aligned}\|u\bar{u}_x\|_{L_x^pL_T^2}=&\left\|\sum_{N_1,N_2}P_{N_1}uP_{N_2}\bar{u}_x\right\|_{L_x^pL_T^2}\\ \lesssim&\left\|\sum_{N_1\sim N_2}P_{N_1}uP_{N_2}\bar{u}_x\right\|_{L_x^pL_T^2}+\left\|\sum_{N_1\ll N_2}P_{N_1}uP_{N_2}\bar{u}_x\right\|_{L_x^pL_T^2} \qquad (4.2)\\ &+\left\|\sum_{N_1\gg N_2}P_{N_1}uP_{N_2}\bar{u}_x\right\|_{L_x^pL_T^2}\\ =:&I_1+I_2+I_3. \qquad (4.3)\end{aligned}$$

Now, we derive the estimates for I_1, I_2 and I_3, respectively.

From the Hölder inequality, the Bernstein type inequalities and the real interpolation theorem, we have

$$\begin{aligned}I_1\lesssim&\sum_{N_1\sim N_2}\|P_{N_1}u\|_{L_x^{2p}L_T^4}\|P_{N_2}u_x\|_{L_x^{2p}L_T^4}\lesssim\sum_{N_1\sim N_2}\|P_{N_1}u\|_{L_x^{2p}L_T^4}N_2\|P_{N_2}u\|_{L_x^{2p}L_T^4}\\ \lesssim&\sum_{N_1\sim N_2}\left\|D_x^{1/2}P_{N_1}u\right\|_{L_x^{2p}L_T^4}\left\|D_x^{1/2}P_{N_2}u\right\|_{L_x^{2p}L_T^4}\lesssim\sum_{N}\left\|D_x^{1/2}P_Nu\right\|_{L_x^{2p}L_T^4}^2\\ \lesssim&\sum_{N}\left\|D_x^{1/2}P_N(P_{\lesssim1}u+P_{\gg1}u)\right\|_{L_x^{2p}L_T^4}^2\\ \lesssim&\sum_{N}\left\|D_x^{1/2}P_NP_{\lesssim1}u\right\|_{L_x^{2p}L_T^4}^2+\sum_{N}\left\|D_x^{1/2}P_NP_{\gg1}u\right\|_{L_x^{2p}L_T^4}^2\\ \lesssim&\sum_{N}\left\|P_NP_{\lesssim1}u\right\|_{L_x^{2p}L_T^4}^2+\sum_{N}N\left\|P_NP_{\gg1}u\right\|_{L_x^{2p}L_T^4}^2\\ \lesssim&\sum_{N\lesssim1}\|P_Nu\|_{L_x^{2p}L_T^4}^2+\sum_{N}N\|P_NP_{\gg1}u\|_{L_x^pL_T^\infty}\|P_NP_{\gg1}u\|_{L_x^\infty L_T^2}.\end{aligned}$$

Applying the Sobolev embedding theorem and the Hölder inequality to the first term, and Bernstein estimates to the second term, we can see that it is bounded by

$$\lesssim T^{1/2}\|u\|_{X_T}^2+\sum_{N}\|P_NP_{\gg1}u\|_{L_x^pL_T^\infty}\|\partial_xP_NP_{\gg1}u\|_{L_x^\infty L_T^2}.$$

By the Cauchy-Schwartz inequality and the Sobolev embedding theorem(i.e. $H_4^{1/4} \subset H_4^{1/4-1/p} \subset L^p$ for the real number $p \geqslant 4$), we can bound it by

$$\lesssim T^{1/2} \|u\|_{X_T}^2 + \|P_{\gg 1} u\|_{X_T}^2 .$$

For I_2 or I_3, it is suffice to consider one of them, e.g. I_2, in view of symmetry. For $N_1 \ll N_2$, we have

$$P_{N_1} u P_{N_2} \bar{u}_x = \tilde{P}_{N_2}(P_{N_1} u P_{N_2} \bar{u}_x),$$

where $\tilde{P}_N = \sum_{j=-2}^{2} P_{2^j N}$. We split these into three cases, i.e. $N_1 \lesssim 1 \ll N_2$, $N_1 \ll N_2 \lesssim 1$ and $1 \ll N_1 \ll N_2$. For the case $N_1 \lesssim 1 \ll N_2$, from the Hölder inequality and the Littlewood-Paley theorem, we can get

$$\begin{aligned}
&\Big\| \sum_{N_1 \lesssim 1 \ll N_2} P_{N_1} u P_{N_2} \bar{u}_x \Big\|_{L_x^p L_T^2} \\
&= \|P_{\lesssim 1} u P_{N_2 \gg 1} \bar{u}_x\|_{L_x^p L_T^2} \lesssim \|P_{\lesssim 1} u\|_{L_x^p L_T^\infty} \|P_{\gg 1} u_x\|_{L_x^\infty L_T^2} \\
&\lesssim \bigg\| \Big(\sum_M |P_M P_{\lesssim 1} u|^2 \Big)^{1/2} \bigg\|_{L_x^p L_T^\infty} \bigg\| \Big(\sum_M |P_M P_{\gg 1} u_x|^2 \Big)^{1/2} \bigg\|_{L_x^\infty L_T^2} \\
&\lesssim \Big(\sum_{M \lesssim 1} \|P_M u\|_{L_x^p L_T^\infty}^2 \Big)^{1/2} \Big(\sum_M \|P_M P_{\gg 1} u_x\|_{L_x^\infty L_T^2}^2 \Big)^{1/2} \\
&\lesssim \|u\|_{X_T} \|P_{\geqslant 1} u\|_{X_T} .
\end{aligned} \tag{4.4}$$

For the case $N_1 \ll N_2 \lesssim 1$, we have, by the Hölder inequality and the Littlewood-Paley theorem, that

$$\begin{aligned}
&\Big\| \sum_{N_1 \ll N_2 \lesssim 1} P_{N_1} u P_{N_2} \bar{u}_x \Big\|_{L_x^p L_T^2} = \Big\| \sum_{N_2 \lesssim 1} P_{\ll N_2} u P_{N_2} \bar{u}_x \Big\|_{L_x^p L_T^2} \\
\lesssim & T^{1/2} \bigg\| \Big(\sum_{N_2 \lesssim 1} |P_{\ll N_2} u|^2 \Big)^{1/2} \bigg\|_{L_x^{2p} L_T^\infty} \bigg\| \Big(\sum_{N_2 \lesssim 1} |P_{N_2} \bar{u}_x|^2 \Big)^{1/2} \bigg\|_{L_x^{2p} L_T^\infty} \\
\lesssim & T^{1/2} \Big(\sum_{N_2 \lesssim 1} \|P_{\ll N_2} u\|_{L_x^{2p} L_T^\infty}^2 \Big)^{1/2} \|u\|_{X_T} .
\end{aligned} \tag{4.5}$$

For $N_2 \lesssim 1$, we have, by the Sobolev embedding theorem, that

$$\|P_{\ll N_2} u\|_{L_x^{2p} L_T^\infty} \sim \bigg\| \Big(\sum_M |P_M P_{\ll N_2} u|^2 \Big)^{1/2} \bigg\|_{L_x^{2p} L_T^\infty} \lesssim \Big(\sum_{M \ll N_2} \|P_M u\|_{L_x^{2p} L_T^\infty}^2 \Big)^{1/2}$$

$$\lesssim \Big(\sum_{M \ll N_2} N_2^{2\varepsilon} M^{-2\varepsilon} \|P_M u\|_{L_x^{2p} L_T^\infty}^2 \Big)^{1/2} \lesssim \Big(\sum_{M \ll N_2} N_2^{2\varepsilon} \|D_x^{-\varepsilon} P_M u\|_{L_x^{2p} L_T^\infty}^2 \Big)^{1/2}$$

$$\lesssim N_2^{\varepsilon}\left(\sum_{M\lesssim 1}\|P_M u\|^2_{L^2_x L^\infty_T}\right)^{1/2}\lesssim N_2^{\varepsilon}\|u\|_{X_T},$$

where $\varepsilon=(p-1)/2p$. Thus, (4.5) can be bounded by

$$\lesssim T^{1/2}\left(\sum_{N_2\lesssim 1}N_2^{2\varepsilon}\right)^{1/2}\|u\|^2_{X_T}\lesssim T^{1/2}\|u\|^2_{X_T}.$$

Now, we turn to the case $1\ll N_1\ll N_2$. From the Littlewood-Paley theorem, we have for $p\geqslant 4$

$$\begin{aligned}\left\|\sum_{1\ll N_1\ll N_2}P_{N_1}uP_{N_2}\bar{u}_x\right\|_{L^p_xL^2_T}&=\left\|\sum_{N_1\gg 1}P_{N_1}uP_{\gg N_1}\bar{u}_x\right\|_{L^p_xL^2_T}\\ \lesssim\sum_{N_1\gg 1}\|P_{N_1}uP_{\gg N_1}\bar{u}_x\|_{L^p_xL^2_T}&\lesssim\sum_{N_1\gg 1}\|P_{N_1}u\|_{L^p_xL^\infty_T}\|P_{\gg N_1}u_x\|_{L^\infty_xL^2_T}.\end{aligned}\tag{4.6}$$

Noticing that

$$\begin{aligned}\|P_{\gg N_1}u_x\|_{L^\infty_xL^2_T}&\sim\left\|\left(\sum_M|P_MP_{\gg N_1}u_x|^2\right)^{1/2}\right\|_{L^\infty_xL^2_T}\lesssim\left(\sum_{M\gg N_1}\|P_Mu_x\|^2_{L^\infty_xL^2_T}\right)^{1/2}\\&\lesssim\|u\|_{X_T},\end{aligned}$$

and for $N_1\gg 1$, $\varepsilon=1/p$ and $p\geqslant 4$

$$\begin{aligned}\|P_{N_1}u\|_{L^p_xL^\infty_T}&=\left\|P_{N_1}(P_{\lesssim 1}u+P_{\gg 1}u)\right\|_{L^p_xL^\infty_T}=\|P_{N_1}P_{\gg 1}u\|_{L^p_xL^\infty_T}\\&=N_1^{-\varepsilon}N_1^{\varepsilon}\|P_{N_1}P_{\gg 1}u\|_{L^p_xL^\infty_T}\sim N_1^{-\varepsilon}\|D_x^{\varepsilon}P_{N_1}P_{\gg 1}u\|_{L^p_xL^\infty_T}\\&\lesssim N_1^{-\varepsilon}\left\|\langle D_x\rangle^{\frac14}P_{N_1}P_{\gg 1}u\right\|_{L^4_xL^\infty_T},\end{aligned}\tag{4.7}$$

we can bound (4.6) by

$$\begin{aligned}&\lesssim\|u\|_{X_T}\sum_{N_1\gg 1}N_1^{-\varepsilon}\left\|\langle D_x\rangle^{\frac14}P_{N_1}P_{\gg 1}u\right\|_{L^4_xL^\infty_T}\\&\lesssim\|u\|_{X_T}\left(\sum_{N_1\gg 1}N_1^{-2\varepsilon}\right)^{\frac12}\left(\sum_{N_1\gg 1}\left\|\langle D_x\rangle^{\frac14}P_{N_1}P_{\gg 1}u\right\|^2_{L^4_xL^\infty_T}\right)^{\frac12}\\&\lesssim\|u\|_{X_T}\|P_{\gg 1}u\|_{X_T},\end{aligned}\tag{4.8}$$

in view of the Hölder inequality. Thus, we have obtained

$$\left\|\sum_{1\ll N_1\ll N_2}P_{N_1}uP_{N_2}\bar{u}_x\right\|_{L^p_xL^2_T}\lesssim\|u\|_{X_T}\|P_{\gg 1}u\|_{X_T},\quad\forall p\geqslant 4.\tag{4.9}$$

Therefore, we have the desired result (4.1) for any real number $p\geqslant 4$. ☐

5. **Nonlinear estimates.** To state the estimates for the nonlinearities $I_{N,j}$, we define the function space Y_T equipped with the following norm:

$$\|u\|_{Y_T}=\|u\|_{L^\infty_TH^{1/2}_x}+\|\partial_xu\|_{L^\infty_xL^2_T}+\|u\|_{L^2_xL^\infty_T}+\left\|\langle D_x\rangle^{\frac14}u\right\|_{L^4_xL^\infty_T}.$$

We have the following proposition for the nonlinearities.

Proposition 5.1. *Let u be a H^∞-solution to* (1.1)-(1.2)*. Then,*

$$
\begin{aligned}
&\left(\sum_{N\gg 1}\left\|\int_0^t S(t-\tau)\sum_{j=1}^{5} I_{N,j}(\tau)d\tau\right\|_{Y_T}^2\right)^{1/2}\\
\lesssim&(1+\|u\|_{X_T}^k)\left[T^{\frac{1}{2}}\|u\|_{X_T}^{\tilde{k}+1}\|P_{\gg 1}u\|_{X_T}^{k-\tilde{k}}+(1+T^{\frac{1}{4}}\|u\|_{X_T})\|u\|_{X_T}^{\tilde{k}}\|P_{\gg 1}u\|_{X_T}^{k+1-\tilde{k}}\right]\\
&+T^{\frac{1}{2}}\left(\|u\|_{X_T}^{2k-1}+\|u\|_{X_T}^{(5k-2)/2}\right)\|P_{\gg 1}u\|_{X_T}\\
&+T^{\frac{1}{4}}(1+T^{\frac{1}{4}}\|u\|_{X_T})^{\frac{1}{2}}\|u\|_{X_T}^{2k-1}\|P_{\gg 1}u\|_{X_T}^{\frac{3}{2}}\\
&+(1+\|u\|_{X_T}^k)\left[T^{\frac{1}{2}}\|u\|_{X_T}^{k}\|P_{\gg 1}u\|_{X_T}+(1+T^{\frac{1}{4}}\|u\|_{X_T})^2\|u\|_{X_T}^{k-1}\|P_{\gg 1}u\|_{X_T}^{2}\right]\\
&+T^{\frac{1}{2}}\|u\|_{X_T}^{3k}\|P_{\gg 1}u\|_{X_T},
\end{aligned}
$$

where $\tilde{k}$ denotes the maximal integer that is less than k (i.e. $\tilde{k}=[k]$ if k is not an integer and $\tilde{k}=k-1$ if k is an integer where $[k]$ denotes the maximal integer that is less than or equal to k).

We consider each nonlinearity separately.

5.1. **Nonlinear estimates of $I_{N,1}$.**

Noting that the term $P_N(|P_{\ll N}u|^k u_x)$ has Fourier support in $|\xi|\sim N$, we have

$$
\begin{aligned}
&P_N(|u|^k u_x)-|P_{\ll N}u|^k P_N u_x\\
=&P_N((|u|^k-|P_{\ll N}u|^k)u_x)+P_N(|P_{\ll N}u|^k u_x)-|P_{\ll N}u|^k P_N u_x\\
=&P_N((|u|^k-|P_{\ll N}u|^k)u_x)+P_N(|P_{\ll N}u|^k \tilde{P}_N u_x)-|P_{\ll N}u|^k P_N\tilde{P}_N u_x, \qquad (5.1)
\end{aligned}
$$

where $\tilde{P}_N=P_{N/2}+P_N+P_{2N}$.

For the second term in (5.1), we have the following estimate.

Lemma 5.1. *Let u be a solution of* (1.1)-(1.2)*. Then, we have for any $k\geqslant 4$*

$$
\begin{aligned}
&\left(\sum_{N\gg 1}\left\|P_N(|P_{\ll N}u|^k\tilde{P}_N u_x)-|P_{\ll N}u|^k P_N\tilde{P}_N u_x\right\|_{L_x^1L_T^2}^2\right)^{1/2} \qquad (5.2)\\
\lesssim&T^{\frac{1}{2}}\|u\|_{X_T}^k\|P_{\gg 1}u\|_{X_T}+(1+T^{\frac{1}{4}}\|u\|_{X_T})\|u\|_{X_T}^{k-1}\|P_{\gg 1}u\|_{X_T}^2.
\end{aligned}
$$

Proof. To shift a derivative from the high-frequency function $P_N u_x$ to the low-frequency function $|P_{\ll N}u|^k$, we require the following Leibniz rule for P_N from [14]:

$$
(P_N(fg)-fP_Ng)(x)=\int_0^1\left(\int\check{\varphi}_N(y)yf_x(x-\eta y)g(x-y)dy\right)d\eta. \qquad (5.3)
$$

Thus, we have

$$
\begin{aligned}
&\left\|P_N(|P_{\ll N}u|^k\tilde{P}_N u_x)-|P_{\ll N}u|^k P_N\tilde{P}_N u_x\right\|_{L_x^1L_T^2}\\
\lesssim&\|\check{\varphi}_N(y)y\|_{L_y^1}\left\|(|P_{\ll N}u|^k)_x\right\|_{L_{x,T}^2}\left\|\tilde{P}_N u_x\right\|_{L_x^2L_T^\infty}\\
\lesssim&N^{-1}\|\check{\varphi}_1(y)y\|_{L_y^1}\left\|(|P_{\ll N}u|^k)_x\right\|_{L_{x,T}^2}\left\|\tilde{P}_N u_x\right\|_{L_x^2L_T^\infty}\\
\lesssim&\left\|(|P_{\ll N}u|^k)_x\right\|_{L_{x,T}^2}\left\|\tilde{P}_N u\right\|_{L_x^2L_T^\infty}
\end{aligned}
$$

$$\lesssim \|P_{\ll N}u\|_{L_x^{2k}L_T^\infty}^{k-2} \left\|P_{\ll N}u_x\overline{P_{\ll N}u}\right\|_{L_x^kL_T^2} \left\|\tilde{P}_Nu\right\|_{L_x^2L_T^\infty}. \tag{5.4}$$

Decomposing $P_{\ll N}u = P_{\leqslant 1}u + P_{1<\cdot\ll N}u$ for $N \gg 1$, we have

$$P_{\ll N}u_x\overline{P_{\ll N}u} = P_{\leqslant 1}u_x\overline{P_{\ll N}u} + P_{1<\cdot\ll N}u_x\overline{P_{\leqslant 1}u} + P_{1<\cdot\ll N}u_x\overline{P_{1<\cdot\ll N}u}. \tag{5.5}$$

For the first term in (5.5), we have

$$\begin{aligned}\left\|P_{\leqslant 1}u_x\overline{P_{\ll N}u}\right\|_{L_x^kL_T^2} \lesssim & \|P_{\leqslant 1}u_x\|_{L_x^{2k}L_T^2} \|P_{\ll N}u\|_{L_x^{2k}L_T^\infty} \\ \lesssim & T^{1/2} \|P_{\leqslant 1}u\|_{L_x^{2k}L_T^\infty} \|P_{\ll N}u\|_{L_x^{2k}L_T^\infty}.\end{aligned}$$

By the Littlewood-Paley theorem, we can obtain

$$\begin{aligned}\|P_{\ll N}u\|_{L_x^{2k}L_T^\infty} \sim & \left\|\left(\sum_M |P_MP_{\ll N}u|^2\right)^{1/2}\right\|_{L_x^{2k}L_T^\infty} \lesssim \left(\sum_M \|P_MP_{\ll N}u\|_{L_x^{2k}L_T^\infty}^2\right)^{1/2} \\ \lesssim & \left(\sum_{M\ll N} \|P_Mu\|_{L_x^{2k}L_T^\infty}^2\right)^{1/2} \lesssim \left(\sum_{M\gg 1} \|P_Mu\|_{L_x^{2k}L_T^\infty}^2\right)^{1/2} \\ \lesssim & \left(\sum_{M\gg 1} \left\|\langle D_x\rangle^{\frac{1}{4}}P_Mu\right\|_{L_x^4L_T^\infty}^2\right)^{1/2} \lesssim \|P_{\gg 1}u\|_{X_T}.\end{aligned} \tag{5.6}$$

In the similar way, we have

$$\|P_{\leqslant 1}u\|_{L_x^{2k}L_T^\infty} \lesssim \|u\|_{X_T}.$$

Thus,

$$\left\|P_{\leqslant 1}u_x\overline{P_{\ll N}u}\right\|_{L_x^kL_T^2} \lesssim T^{1/2} \|u\|_{X_T} \|P_{\gg 1}u\|_{X_T}. \tag{5.7}$$

For the last two term in (5.5), in a similar way as in the proof of Proposition 4.1, we can obtain the following bound:

$$\|P_{\leqslant 1}\bar{u}P_{1\ll\cdot\ll N}u_x\|_{L_x^kL_T^2} \lesssim T^{\frac{1}{2}} \|u\|_{X_T}^2 + \|u\|_{X_T} \|P_{\gg 1}u\|_{X_T}, \tag{5.8}$$

$$\|P_{1\ll\cdot\ll N}\bar{u}P_{1\ll\cdot\ll N}u_x\|_{L_x^kL_T^2} \lesssim T^{\frac{1}{2}} \|u\|_{X_T}^2 + (1+T^{\frac{1}{4}} \|u\|_{X_T}) \|u\|_{X_T} \|P_{\gg 1}u\|_{X_T}. \tag{5.9}$$

From the Sobolev embedding theorem and (5.7)-(5.9), we obtain that (5.4) can be bounded by

$$\lesssim \left(T^{\frac{1}{2}} \|u\|_{X_T}^k + (1+T^{\frac{1}{4}} \|u\|_{X_T}) \|u\|_{X_T}^{k-1} \|P_{\gg 1}u\|_{X_T}\right) \left\|\tilde{P}_Nu\right\|_{L_x^2L_T^\infty}.$$

Thus, we can bound (5.2) by

$$\begin{aligned}&\lesssim \left(T^{\frac{1}{2}} \|u\|_{X_T}^k + (1+T^{\frac{1}{4}} \|u\|_{X_T}) \|u\|_{X_T}^{k-1} \|P_{\gg 1}u\|_{X_T}\right) \left(\sum_{N\gg 1} \|P_Nu\|_{L_x^2L_T^\infty}^2\right)^{1/2} \\ &\lesssim T^{\frac{1}{2}} \|u\|_{X_T}^k \|P_{\gg 1}u\|_{X_T} + (1+T^{\frac{1}{4}} \|u\|_{X_T}) \|u\|_{X_T}^{k-1} \|P_{\gg 1}u\|_{X_T}^2,\end{aligned}$$

which yields the desired result. □

For the first term in (5.1), we have the following estimate:

Lemma 5.2. *Let u be a solution of* (1.1)-(1.2). *Then, we have for any $k \geqslant 4$*

$$\begin{aligned}&\left(\sum_{N\gg 1}\left\|P_N((|u|^k-|P_{\ll N}u|^k)u_x)\right\|^2_{L^1_xL^2_T}\right)^{1/2}\\ \lesssim & T^{\frac{1}{2}}\left\|u\right\|^{\tilde{k}+1}_{X_T}\left\|P_{\gg 1}u\right\|^{k-\tilde{k}}_{X_T}+(1+T^{\frac{1}{4}}\left\|u\right\|_{X_T})\left\|u\right\|^{\tilde{k}}_{X_T}\left\|P_{\gg 1}u\right\|^{k+1-\tilde{k}}_{X_T}. \qquad (5.10)\end{aligned}$$

Proof. We split (5.1) into several terms for $N \gg 1$ and $k \geqslant 4$

$$\begin{aligned}&P_N((|u|^k-|P_{\ll N}u|^k)u_x) &(5.11)\\ =&P_N(|u|^{k-2}\bar{u}u_xP_{\gtrsim N}u) &(5.12)\\ &+P_N((|u|^{k-2}-|P_{\ll N}u|^{k-2})\bar{u}u_xP_{\ll N}u) &(5.13)\\ &+P_N(|P_{\ll N}u|^{k-2}u_xP_{\ll N}uP_{\gtrsim N}\bar{u}). &(5.14)\end{aligned}$$

Notice that

$$\begin{aligned}\left\|P_{\gtrsim N}u\right\|_{L^k_xL^\infty_T}\lesssim&\left(\sum_{M\gtrsim N}\left\|P_Mu\right\|^2_{L^k_xL^\infty_T}\right)^{1/2}\lesssim\left(\sum_{M\gtrsim N}N^{-2\varepsilon_k}M^{2\varepsilon_k}\left\|P_Mu\right\|^2_{L^k_xL^\infty_T}\right)^{1/2}\\ \lesssim&\left(\sum_{M\gtrsim N}N^{-2\varepsilon_k}\left\|D^{\varepsilon_k}_xP_Mu\right\|^2_{L^k_xL^\infty_T}\right)^{1/2}\\ \lesssim&N^{-\varepsilon_k}\left(\sum_{M\gtrsim N}\left\|\langle D_x\rangle^{\frac{1}{4}}P_Mu\right\|^2_{L^4_xL^\infty_T}\right)^{1/2}\\ \lesssim&N^{-\varepsilon_k}\left\|P_{\gg 1}u\right\|_{X_T},\quad\forall k\geqslant 4,\end{aligned}$$

where $\varepsilon_k > 0$ is defined by $\varepsilon_k = 1/k$.

Thus, for the first term (5.12), from the fact $\|\check{\varphi}_N\|_{L^1} \lesssim 1$ and Proposition 4.1, we have for $k \geqslant 4$

$$\begin{aligned}&\left\|P_N(|u|^{k-2}\bar{u}u_xP_{\gtrsim N}u)\right\|_{L^1_xL^2_T}\lesssim\left\||u|^{k-2}\bar{u}u_xP_{\gtrsim N}u\right\|_{L^1_xL^2_T}\\ \lesssim&\left\|u\right\|^{k-2}_{L^k_xL^\infty_T}\left\|\bar{u}u_x\right\|_{L^k_xL^2_T}\left\|P_{\gtrsim N}u\right\|_{L^k_xL^\infty_T}\\ \lesssim&N^{-\varepsilon_k}\left[T^{\frac{1}{2}}\left\|u\right\|^k_{X_T}\left\|P_{\gg 1}u\right\|_{X_T}+(1+T^{\frac{1}{4}}\left\|u\right\|_{X_T})\left\|u\right\|^k_{X_T}\left\|P_{\gg 1}u\right\|^2_{X_T}\right].\end{aligned}$$

Therefore, we obtain, for any $k \geqslant 4$, that

$$\begin{aligned}&\left(\sum_{N\gg 1}\left\|P_N(|u|^{k-2}\bar{u}u_xP_{\gtrsim N}u)\right\|^2_{L^1_xL^2_T}\right)^{1/2}\\ \lesssim&T^{\frac{1}{2}}\left\|u\right\|^k_{X_T}\left\|P_{\gg 1}u\right\|_{X_T}+(1+T^{\frac{1}{4}}\left\|u\right\|_{X_T})\left\|u\right\|^k_{X_T}\left\|P_{\gg 1}u\right\|^2_{X_T}.\end{aligned}$$

For (5.14), in the same way as the case (5.12), we have

$$\begin{aligned}&\left(\sum_{N\gg 1}\left\|P_N(|P_{\ll N}u|^{k-2}u_xP_{\ll N}uP_{\gtrsim N}\bar{u})\right\|^2_{L^1_xL^2_T}\right)^{1/2}\\ \lesssim&T^{\frac{1}{2}}\left\|u\right\|^k_{X_T}\left\|P_{\gg 1}u\right\|_{X_T}+(1+T^{\frac{1}{4}}\left\|u\right\|_{X_T})\left\|u\right\|^k_{X_T}\left\|P_{\gg 1}u\right\|^2_{X_T}.\end{aligned}$$

Now, we derive the estimate for (5.13) by using the induction argument in k.

For $k=4$, we have

$$\left||u|^{k-2}-|P_{\ll N}u|^{k-2}\right|\lesssim\left|P_{\gtrsim N}u\right|^2+\left|P_{\gtrsim N}u\overline{P_{\ll N}u}\right|.$$

From the Young inequality, the Hölder inequality, (5.6) and Proposition 4.1, we can get for $k=4$

$$\begin{aligned}
&\left\|P_N((|u|^{k-2}-|P_{\ll N}u|^{k-2})\bar{u}u_xP_{\ll N}u)\right\|_{L^1_xL^2_T}\\
\lesssim&\left\||u|^{k-2}-|P_{\ll N}u|^{k-2}\right\|_{L_x^{2k/(k-2)}L_T^\infty}\left\|\bar{u}u_x\right\|_{L^k_xL^2_T}\left\|P_{\ll N}u\right\|_{L^2_xL^\infty_T}\\
\lesssim&(\left\|P_{\gtrsim N}u\right\|^2_{L^8_xL^\infty_T}+\left\|P_{\gtrsim N}u\right\|_{L^8_xL^\infty_T}\left\|P_{\ll N}u\right\|_{L^8_xL^\infty_T})\left\|\bar{u}u_x\right\|_{L^k_xL^2_T}\left\|P_{\ll N}u\right\|_{L^2_xL^\infty_T}\\
\lesssim&N^{-1/8}\left\|P_{\gg1}u\right\|^2_{X_T}\left\|\bar{u}u_x\right\|_{L^k_xL^2_T}\left\|P_{\ll N}u\right\|_{L^2_xL^\infty_T}\\
\lesssim&N^{-1/8}\left[T^{\frac12}\left\|u\right\|^3_{X_T}\left\|P_{\gg1}u\right\|^{k-2}_{X_T}+(1+T^{\frac14}\left\|u\right\|_{X_T})\left\|u\right\|^2_{X_T}\left\|P_{\gg1}u\right\|^{k-1}_{X_T}\right].
\end{aligned}$$

From the triangle inequality for complex number, i.e. $||z_1|-|z_2||\leqslant|z_1-z_2|$ for $z_1,z_2\in\mathbb{C}$, we can get $\left||z_1|^\theta-|z_2|^\theta\right|\leqslant|z_1-z_2|^\theta$ for any $\theta\in(0,1]$.

For $k\in(4,5]$, we have

$$\begin{aligned}
&\left||u|^{k-2}-|P_{\ll N}u|^{k-2}\right|\\
\lesssim&|u|^2\left||u|^{k-4}-|P_{\ll N}u|^{k-4}\right|+|P_{\ll N}u|^{k-4}\left||u|^2-|P_{\ll N}u|^2\right|\\
\lesssim&|u|^2\left|P_{\gtrsim N}u\right|^{k-4}+|P_{\ll N}u|^{k-4}\left|P_{\gtrsim N}u\right|^2+\left|P_{\gtrsim N}u\right|\left|P_{\ll N}u\right|^{k-3}.
\end{aligned}$$

Then

$$\begin{aligned}
&\left\|P_N((|u|^{k-2}-|P_{\ll N}u|^{k-2})\bar{u}u_xP_{\ll N}u)\right\|_{L^1_xL^2_T}\\
\lesssim&\Big[\left\|u\right\|^2_{L^{2k}_xL^\infty_T}\left\|P_{\gtrsim N}u\right\|^{k-4}_{L^{2k}_xL^\infty_T}+\left\|P_{\ll N}u\right\|^{k-4}_{L^{2k}_xL^\infty_T}\left\|P_{\gtrsim N}u\right\|^2_{L^{2k}_xL^\infty_T}\\
&\quad+\left\|P_{\gtrsim N}u\right\|_{L^{2k}_xL^\infty_T}\left\|P_{\ll N}u\right\|^{k-3}_{L^{2k}_xL^\infty_T}\Big]\left\|\bar{u}u_x\right\|_{L^k_xL^2_T}\left\|P_{\ll N}u\right\|_{L^2_xL^\infty_T}\\
\lesssim&N^{-\varepsilon_k}\left[T^{\frac12}\left\|u\right\|^4_{X_T}\left\|P_{\gg1}u\right\|^{k-3}_{X_T}+(1+T^{\frac14}\left\|u\right\|_{X_T})\left\|u\right\|^3_{X_T}\left\|P_{\gg1}u\right\|^{k-2}_{X_T}\right],
\end{aligned}$$

where $\varepsilon_k=(k-4)/2k$ for $k\in(4,5]$. By the same procedure, we can obtain for any $k\geqslant4$

$$\begin{aligned}
&\left\|P_N((|u|^{k-2}-|P_{\ll N}u|^{k-2})\bar{u}u_xP_{\ll N}u)\right\|_{L^1_xL^2_T}\\
\lesssim&N^{-\varepsilon_k}\left[T^{\frac12}\left\|u\right\|^{\tilde{k}+1}_{X_T}\left\|P_{\gg1}u\right\|^{k-\tilde{k}}_{X_T}+(1+T^{\frac14}\left\|u\right\|_{X_T})\left\|u\right\|^{\tilde{k}}_{X_T}\left\|P_{\gg1}u\right\|^{k+1-\tilde{k}}_{X_T}\right],
\end{aligned}$$

where $\varepsilon_k=(k-\tilde{k})/2k>0$. Therefore, we have for any $k\geqslant4$

$$\begin{aligned}
&\left(\sum_{N\gg1}\left\|P_N((|u|^{k-2}-|P_{\ll N}u|^{k-2})\bar{u}u_xP_{\ll N}u)\right\|^2_{L^1_xL^2_T}\right)^{1/2}\\
\lesssim&T^{\frac12}\left\|u\right\|^{\tilde{k}+1}_{X_T}\left\|P_{\gg1}u\right\|^{k-\tilde{k}}_{X_T}+(1+T^{\frac14}\left\|u\right\|_{X_T})\left\|u\right\|^{\tilde{k}}_{X_T}\left\|P_{\gg1}u\right\|^{k+1-\tilde{k}}_{X_T}.
\end{aligned}\tag{5.15}$$

Thus, we have proved the desired result. □

Remark 5.1. From the proof of Lemma 5.2, we can see that

$$\left(\sum_{N\gg1}\left\|P_N((|u|^k-|P_{\ll N}u|^k)u_x)\right\|^2_{L_x^{\frac{1}{1-\varepsilon}}L^2_T}\right)^{1/2}$$

$$\lesssim T^{\frac{1}{2}} \|u\|_{X_T}^{\tilde{k}+1} \|P_{\gg 1}u\|_{X_T}^{k-\tilde{k}} + (1+T^{\frac{1}{4}} \|u\|_{X_T}) \|u\|_{X_T}^{\tilde{k}} \|P_{\gg 1}u\|_{X_T}^{k+1-\tilde{k}}, \tag{5.16}$$

holds for any $\varepsilon \in [0,1)$ in view of Proposition 4.1.

We turn to the proof of Proposition 5.1 for the nonlinearity $I_{N,1}$. We also consider the decomposition in (5.1). For convenience, we denote $B_N = P_N(|P_{\ll N}u|^k \tilde{P}_N u_x) - |P_{\ll N}u|^k P_N \tilde{P}_N u_x$. From (3.5), (3.11), (3.6) and (3.7), we have

$$\begin{aligned} &\left(\sum_{N\gg 1} \left\| \int_0^t S(t-\tau) e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy} B_N d\tau \right\|_{Y_T}^2 \right)^{1/2} \\ &\lesssim \left(\sum_{N\gg 1} \|B_N\|_{L_x^1 L_T^2}^2 \right)^{1/2} + \left(\sum_{N\gg 1} \left(\sum_M \left\| P_M(e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy} B_N) \right\|_{L_x^1 L_T^2} \right)^2 \right)^{1/2}. \end{aligned} \tag{5.17}$$

By Lemma 5.1, the first term can be bounded by

$$\lesssim T^{\frac{1}{2}} \|u\|_{X_T}^k \|P_{\gg 1}u\|_{X_T} + (1+T^{\frac{1}{4}} \|u\|_{X_T}) \|u\|_{X_T}^{k-1} \|P_{\gg 1}u\|_{X_T}^2 .$$

For the second term, we split the sum $\sum_M$ into three parts $\sum_{M\sim N} + \sum_{M\ll N} + \sum_{M\gg N}$ as in [14]. For the part of $M \sim N$, it is the same as Lemma 5.1 by summing in M such that $M \sim N$. For the part $M \ll N$, we can add the projection operator $P_{\sim N}$ to $e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy}$ since B_N has Fourier support in $|\xi| \sim N$. Thus, by the Hölder inequality, we have

$$\begin{aligned} &\left(\sum_{N\gg 1} \left(\sum_{M\ll N} \left\| P_M(e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy} B_N) \right\|_{L_x^1 L_T^2} \right)^2 \right)^{1/2} \\ &\lesssim \left(\sum_{N\gg 1} \left(\sum_{M\ll N} \left\| P_{\sim N} e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy} B_N \right\|_{L_x^1 L_T^2} \right)^2 \right)^{1/2} \\ &\lesssim \left(\sum_{N\gg 1} (\ln N)^2 \left\| P_{\sim N} e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy} \right\|_{L_x^{1/\varepsilon} L_T^\infty}^2 \|B_N\|_{L_x^{1/(1-\varepsilon)} L_T^2}^2 \right)^{1/2}, \end{aligned} \tag{5.18}$$

where $\varepsilon \in (0, 1/k)$.

By the Bernstein inequality, we have

$$\begin{aligned} &N \left\| P_{\sim N} e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy} \right\|_{L_x^{1/\varepsilon} L_T^\infty} \lesssim \left\| \partial_x P_{\sim N} e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy} \right\|_{L_x^{1/\varepsilon} L_T^\infty} \\ &\lesssim \|P_{\ll N}u\|_{L_x^{k/\varepsilon} L_T^\infty}^k \lesssim \|u\|_{X_T}^k , \end{aligned}$$

and from (5.3) and the Hölder inequality, we can get, as a similar way as in (5.4), that

$$\begin{aligned} &\|B_N\|_{L_x^{1/(1-\varepsilon)} L_T^2} = \left\| (|P_{\ll N}u|^k)_x \right\|_{L_x^{2/(1-2\varepsilon)} L_T^2} \left\| \tilde{P}_N u \right\|_{L_x^2 L_T^\infty} \\ &\lesssim \|P_{\ll N}u\|_{L_x^{\frac{2k(k-2)}{k-2-2k\varepsilon}} L_T^\infty}^{k-2} \|P_{\ll N}u_x P_{\ll N}\bar{u}\|_{L_x^k L_T^2} \left\| \tilde{P}_N u \right\|_{L_x^2 L_T^\infty} \\ &\lesssim \left(T^{\frac{1}{2}} \|u\|_{X_T}^k + (1+T^{\frac{1}{4}} \|u\|_{X_T}) \|u\|_{X_T}^{k-1} \|P_{\gg 1}u\|_{X_T} \right) \|P_N u\|_{L_x^2 L_T^\infty} . \end{aligned}$$

Thus, (5.18) can be bounded by

$$\begin{aligned}&\lesssim \|u\|_{X_T}^{k}\left(T^{\frac{1}{2}}\|u\|_{X_T}^{k}+(1+T^{\frac{1}{4}}\|u\|_{X_T})\|u\|_{X_T}^{k-1}\|P_{\gg 1}u\|_{X_T}\right)\left(\sum_{N\gg 1}\|P_N u\|^2_{L^2_x L^\infty_T}\right)^{1/2}\\&\lesssim T^{\frac{1}{2}}\|u\|_{X_T}^{2k}\|P_{\gg 1}u\|_{X_T}+(1+T^{\frac{1}{4}}\|u\|_{X_T})\|u\|_{X_T}^{2k-1}\|P_{\gg 1}u\|^2_{X_T}.\end{aligned}$$

For the part $M \gg N$, we can add the projection operator P_M to $e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u|^k dy}$. In a similar way with the part $M \ll N$, we have

$$\begin{aligned}&\left(\sum_{N\gg 1}\left(\sum_{M\gg N}\left\|P_M(e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u|^k dy}B_N)\right\|_{L^1_x L^2_T}\right)^2\right)^{1/2}\\&\lesssim\left(\sum_{N\gg 1}\left(\sum_{M\gg N}\left\|P_M e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u|^k dy}\right\|_{L^{1/\varepsilon}_x L^\infty_T}\|B_N\|_{L^{1/(1-\varepsilon)}_x L^2_T}\right)^2\right)^{1/2}\\&\lesssim\Bigg(\sum_{N\gg 1}\Big(\sum_{M\gg N}\frac{1}{M}\Big(T^{\frac{1}{2}}\|u\|_{X_T}^{2k}\|P_N u\|_{L^2_x L^\infty_T}\\&\qquad\qquad+(1+T^{\frac{1}{4}}\|u\|_{X_T})\|u\|_{X_T}^{2k-1}\|P_{\gg 1}u\|_{X_T}\|P_N u\|_{L^2_x L^\infty_T}\Big)\Big)^2\Bigg)^{1/2}\\&\lesssim T^{\frac{1}{2}}\|u\|_{X_T}^{2k}\|P_{\gg 1}u\|_{X_T}+(1+T^{\frac{1}{4}}\|u\|_{X_T})\|u\|_{X_T}^{2k-1}\|P_{\gg 1}u\|^2_{X_T}.\end{aligned}$$

For the first term in (5.1), we denote it by A_N, i.e. $A_N = P_N((|u|^k - |P_{\ll N}u|^k)u_x)$. Similarly, from (3.5), (3.11), (3.6) and (3.7), we can get

$$\begin{aligned}&\left(\sum_{N\gg 1}\left\|\int_0^t S(t-\tau)e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u|^k dy}A_N d\tau\right\|^2_{Y_T}\right)^{1/2}\\&\lesssim\left(\sum_{N\gg 1}\|A_N\|^2_{L^1_x L^2_T}\right)^{1/2}+\left(\sum_{N\gg 1}\left(\sum_{M}\left\|P_M(e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u|^k dy}A_N)\right\|_{L^1_x L^2_T}\right)^2\right)^{1/2}.\end{aligned}\tag{5.19}$$

From Lemma 5.2, the first term is bounded by

$$\lesssim T^{\frac{1}{2}}\|u\|_{X_T}^{\tilde{k}+1}\|P_{\gg 1}u\|_{X_T}^{k-\tilde{k}}+(1+T^{\frac{1}{4}}\|u\|_{X_T})\|u\|_{X_T}^{\tilde{k}}\|P_{\gg 1}u\|_{X_T}^{k+1-\tilde{k}}.$$

Noticing that (5.16), and in the same way as in dealing with the second term of (5.17), we can bound the second term of (5.19) by

$$\lesssim T^{\frac{1}{2}}\|u\|_{X_T}^{k+\tilde{k}+1}\|P_{\gg 1}u\|_{X_T}^{k-\tilde{k}}+(1+T^{\frac{1}{4}}\|u\|_{X_T})\|u\|_{X_T}^{k+\tilde{k}}\|P_{\gg 1}u\|_{X_T}^{k+1-\tilde{k}}.$$

Therefore, we have obtained

$$\begin{aligned}&\left(\sum_{N\gg 1}\left\|\int_0^t S(t-\tau)I_{N,1}(\tau)d\tau\right\|^2_{Y_T}\right)^{1/2}\\&\lesssim(1+\|u\|_{X_T}^{k})\left[T^{\frac{1}{2}}\|u\|_{X_T}^{\tilde{k}+1}\|P_{\gg 1}u\|_{X_T}^{k-\tilde{k}}+(1+T^{\frac{1}{4}}\|u\|_{X_T})\|u\|_{X_T}^{\tilde{k}}\|P_{\gg 1}u\|_{X_T}^{k+1-\tilde{k}}\right].\end{aligned}$$

5.2. **Nonlinear estimates of $I_{N,2}$.** From (3.4), (3.8), (3.9) and (3.10), we have

$$\begin{aligned}
&\left(\sum_{N\gg 1}\left\|\int_0^t S(t-\tau)I_{N,2}(\tau)d\tau\right\|_{Y_T}^2\right)^{1/2}\\
\lesssim&\left(\sum_{N\gg 1}\left\|e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy}P_N u B_{N,2}\right\|_{L_T^1 H_x^{1/2}}^2\right)^{1/2} &(5.20)\\
&+\left(\sum_{N\gg 1}\left(\sum_M \left\|P_M(e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy}P_N u B_{N,2})\right\|_{L_T^1 H_x^{1/2}}\right)^2\right)^{1/2}, &(5.21)
\end{aligned}$$

where $B_{N,2}=\int_{-\infty}^x |P_{\ll N}u|^{k-4}\left[(\overline{P_{\ll N}u_x}P_{\ll N}u)^2-(P_{\ll N}u_x\overline{P_{\ll N}u})^2\right]dy$.

For the first term (5.20), from Lemma 3.3 and the Hölder inequality, it can be bounded by

$$\begin{aligned}
\lesssim&\left(\sum_{N\gg 1}\|P_N u B_{N,2}\|_{L_T^1 L_x^2}^2\right.\\
&\qquad\left.+\|P_N u B_{N,2}\|_{L_T^1 L_x^2}\left\|\partial_x(e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy}P_N u B_{N,2})\right\|_{L_T^1 L_x^2}\right)^{1/2}\\
\lesssim&\Big(\sum_{N\gg 1}\|P_N u\|_{L_T^\infty L_x^2}^2\|B_{N,2}\|_{L_T^1 L_x^\infty}^2+\|P_N u\|_{L_T^\infty H_x^{1/2}}^2\|B_{N,2}\|_{L_T^1 L_x^\infty}^2\\
&+\|P_N u\|_{L_T^\infty L_x^2}\|B_{N,2}\|_{L_T^1 L_x^\infty}^2\|P_{\ll N}u\|_{L_x^{2(k+1)}L_T^\infty}^k\|P_N u\|_{L_x^{2(k+1)}L_T^\infty}\\
&+\|P_N u\|_{L_T^\infty L_x^2}\|B_{N,2}\|_{L_T^1 L_x^\infty}\|P_N u\|_{L_x^\infty L_T^2}\|\partial_x B_{N,2}\|_{L_{x,T}^2}\Big)^{1/2}. &(5.22)
\end{aligned}$$

By the Hölder inequality, we have for $k\geqslant 5$

$$\begin{aligned}
\|B_{N,2}\|_{L_T^1 L_x^\infty}\lesssim&\left\||P_{\ll N}u|^{k-4}\left[(\overline{P_{\ll N}u_x}P_{\ll N}u)^2-(P_{\ll N}u_x\overline{P_{\ll N}u})^2\right]\right\|_{L_{x,T}^1}\\
\lesssim&\left\|\overline{P_{\ll N}u}P_{\ll N}u_x\right\|_{L_x^4 L_T^2}^2\|P_{\ll N}u\|_{L_x^{2(k-4)}L_T^\infty}^{k-4}\\
\lesssim&T\|u\|_{X_T}^k+(1+T^{\frac14}\|u\|_{X_T})^2\|u\|_{X_T}^{k-2}\|P_{\gg 1}u\|_{X_T}^2, &(5.23)
\end{aligned}$$

and from Proposition 4.1 and the proof of Lemma 5.1,

$$\begin{aligned}
&\|P_N u\|_{L_x^\infty L_T^2}\|\partial_x B_{N,2}\|_{L_{x,T}^2}\\
=&\|P_N u\|_{L_x^\infty L_T^2}\left\||P_{\ll N}u|^{k-4}\left[(\overline{P_{\ll N}u_x}P_{\ll N}u)^2-(P_{\ll N}u_x\overline{P_{\ll N}u})^2\right]\right\|_{L_{x,T}^2}\\
\lesssim&\|P_N u_x\|_{L_x^\infty L_T^2}\|P_{\ll N}u_x P_{\ll N}\bar{u}\|_{L_x^{2(k-1)}L_T^2}\|P_{\ll N}u\|_{L_x^{2(k-1)}L_T^\infty}^{k-2}\\
\lesssim&\|P_N u_x\|_{L_x^\infty L_T^2}\left[T^{\frac12}\|u\|_{X_T}^k+(1+T^{\frac14}\|u\|_{X_T})\|u\|_{X_T}^{k-1}\|P_{\gg 1}u\|_{X_T}\right].
\end{aligned}$$

Thus, we can bound (5.20) by

$$\lesssim(1+\|u\|_{X_T}^k)\left(T^{1/2}\|u\|_{X_T}^k\|P_{\gg 1}u\|_{X_T}+(1+T^{\frac14}\|u\|_{X_T})^2\|u\|_{X_T}^{k-2}\|P_{\gg 1}u\|_{X_T}^3\right).$$

For (5.21), we split the sum $\sum_M$ into two parts $\sum_{M\lesssim N}+\sum_{M\gg N}$, which gives the bound by

$$\lesssim \left(\sum_{N\gg 1}\left(\sum_{M\lesssim N}\langle M\rangle^{\frac12}\left\|P_NuB_{N,2}\right\|_{L_T^1L_x^2}\right)^2\right)^{1/2} \tag{5.24}$$

$$+\left(\sum_{N\gg 1}\left(\sum_{M\gg N}\left\|P_M\langle D_x\rangle^{\frac12}e^{-\frac{i\lambda}{2}\int_{-\infty}^x|P_{\ll N}u|^k dy}P_NuB_{N,2}\right\|_{L_T^1L_x^2}\right)^2\right)^{1/2}. \tag{5.25}$$

For the first term (5.24), noticing that $\sum_{M\lesssim N}\langle M\rangle^{1/2}\lesssim N^{1/2}$ and (5.23), we can bound it by

$$\begin{aligned}&\lesssim\left(\sum_{N\gg 1}\left(\left\|D_x^{\frac12}P_Nu\right\|_{L_T^\infty L_x^2}\left\|B_{N,2}\right\|_{L_T^1L_x^\infty}\right)^2\right)^{1/2}\\ &\lesssim\|u\|_{X_T}^k\left\|P_{\gg 1}u\right\|_{X_T}.\end{aligned}$$

For the second term (5.25), in a similar way with (5.22), we bound it by

$$\begin{aligned}&\lesssim\left(\sum_{N\gg 1}\left(\sum_{M\gg N}M^{-\frac12}\left\|P_M\langle D_x\rangle e^{-\frac{i\lambda}{2}\int_{-\infty}^x|P_{\ll N}u|^k dy}P_NuB_{N,2}\right\|_{L_T^1L_x^2}\right)^2\right)^{1/2}\\ &\lesssim\left(\sum_{N\gg 1}\left\|P_NuB_{N,2}\right\|_{L_T^1L_x^2}^2+\left\|\partial_x e^{-\frac{i\lambda}{2}\int_{-\infty}^x|P_{\ll N}u|^k dy}P_NuB_{N,2}\right\|_{L_T^1L_x^2}^2\right)^{1/2}\\ &\lesssim(1+\|u\|_{X_T}^k)\left(T^{1/2}\|u\|_{X_T}^k\left\|P_{\gg 1}u\right\|_{X_T}+(1+T^{\frac14}\|u\|_{X_T})^2\|u\|_{X_T}^{k-2}\left\|P_{\gg 1}u\right\|_{X_T}^3\right).\end{aligned}$$

Therefore, we obtain

$$\begin{aligned}&\left(\sum_{N\gg 1}\left\|\int_0^t S(t-\tau)I_{N,2}(\tau)d\tau\right\|_{Y_T}^2\right)^{1/2}\\ &\lesssim(1+\|u\|_{X_T}^k)\left(T^{1/2}\|u\|_{X_T}^k\left\|P_{\gg 1}u\right\|_{X_T}+(1+T^{\frac14}\|u\|_{X_T})^2\|u\|_{X_T}^{k-2}\left\|P_{\gg 1}u\right\|_{X_T}^3\right).\end{aligned}$$

5.3. **Nonlinear estimates of $I_{N,3}$.** From (3.4), (3.8), (3.9) and (3.10), we have

$$\left(\sum_{N\gg 1}\left\|\int_0^t S(t-\tau)I_{N,2}(\tau)d\tau\right\|_{Y_T}^2\right)^{1/2} \tag{5.26}$$

$$\lesssim\left(\sum_{N\gg 1}\left\|e^{-\frac{i\lambda}{2}\int_{-\infty}^x|P_{\ll N}u|^k dy}P_NuB_{N,3}\right\|_{L_T^1H_x^{1/2}}^2\right)^{1/2} \tag{5.27}$$

$$+\left(\sum_{N\gg 1}\left(\sum_{M}\left\|P_M(e^{-\frac{i\lambda}{2}\int_{-\infty}^x|P_{\ll N}u|^k dy}P_NuB_{N,3})\right\|_{L_T^1H_x^{1/2}}\right)^2\right)^{1/2}, \tag{5.28}$$

where $B_{N,3}=\int_{-\infty}^x|P_{\ll N}u|^{k-2}P_{\ll N}|u|^k(u_x+\bar{u}_x)dy$.

By Hölder inequality, we get

$$\left\|B_{N,3}\right\|_{L_T^1L_x^\infty}\lesssim\left\||P_{\ll N}u|^{k-2}P_{\ll N}|u|^k(u_x+\bar{u}_x)\right\|_{L_{x,T}^1}$$

$$
\begin{aligned}
&\lesssim T^{\frac{1}{2}} \left\|P_{\ll N} u\right\|_{L_x^{2k-2} L_T^{\infty}}^{k-2} \left\|P_{\ll N}|u|^k(u_x+\bar{u}_x)\right\|_{L_x^{(2k-2)/k} L_T^2} \\
&\lesssim T^{\frac{1}{2}} \left\|P_{\ll N} u\right\|_{L_x^{2k-2} L_T^{\infty}}^{k-2} \|u\|_{L_x^{2k-2} L_T^{\infty}}^{k} \|u_x\|_{L_x^{\infty} L_T^2} \\
&\lesssim T^{\frac{1}{2}} \|u\|_{X_T}^{2k-1}.
\end{aligned}
$$

From the Hölder inequality and Proposition 4.1, we have

$$
\begin{aligned}
\|\partial_x B_{N,3}\|_{L_{x,T}^2} &= \left\| \left|P_{\ll N} u\right|^{k-2} P_{\ll N}|u|^k(u_x+\bar{u}_x)\right\|_{L_{x,T}^2} \\
&\lesssim \left\|P_{\ll N} u\right\|_{L_x^{4k-4} L_T^{\infty}}^{k-2} \|u\|_{L_x^{4k-4} L_T^{\infty}}^{k} \|\bar{u} u_x\|_{L_x^{4k-4} L_T^2} \\
&\lesssim T^{\frac{1}{2}} \|u\|_{X_T}^{2k} + (1+T^{\frac{1}{4}} \|u\|_{X_T}) \|u\|_{X_T}^{2k-1} \|P_{\gg 1} u\|_{X_T}.
\end{aligned}
$$

In addition, for $N \gg 1$, we have $\|P_N u\|_{L_x^{\infty} L_T^2} \lesssim \|P_N u_x\|_{L_x^{\infty} L_T^2}$. Thus, in the same way as in the case $I_{N,2}$, we can bound (5.26) by

$$
\begin{aligned}
&\lesssim T^{\frac{1}{2}} \left(\|u\|_{X_T}^{2k-1} + \|u\|_{X_T}^{(5k-2)/2} \right) \|P_{\gg 1} u\|_{X_T} \\
&\quad + T^{\frac{1}{4}} (1+T^{\frac{1}{4}} \|u\|_{X_T})^{\frac{1}{2}} \|u\|_{X_T}^{2k-1} \|P_{\gg 1} u\|_{X_T}^{\frac{3}{2}}.
\end{aligned}
$$

5.4. **Nonlinear estimates of $I_{N,4}$.** From (3.5), (3.11), (3.6) and (3.7), we have

$$
\begin{aligned}
&\left(\sum_{N \gg 1} \left\| \int_0^t S(t-\tau) e^{-\frac{i\lambda}{2} \int_{-\infty}^x |P_{\ll N} u|^k dy} B_{N,4} d\tau \right\|_{Y_T}^2 \right)^{1/2} \\
&\lesssim \left(\sum_{N \gg 1} \|B_{N,4}\|_{L_x^1 L_T^2}^2 \right)^{1/2} \\
&\quad + \left(\sum_{N \gg 1} \left(\sum_{M} \left\| P_M (e^{-\frac{i\lambda}{2} \int_{-\infty}^x |P_{\ll N} u|^k dy} B_{N,4}) \right\|_{L_x^1 L_T^2} \right)^2 \right)^{1/2},
\end{aligned} \tag{5.29}
$$

where $B_{N,4} = |P_{\ll N} u|^{k-2} P_N u P_{\ll N} u \overline{P_{\ll N} u_x}$. By the Hölder inequality, we have

$$
\begin{aligned}
\|B_{N,4}\|_{L_x^1 L_T^2} &\lesssim \|P_{\ll N} u\|_{L_x^k L_T^{\infty}}^{k-2} \|P_N u\|_{L_x^k L_T^{\infty}} \left\| P_{\ll N} u \overline{P_{\ll N} u_x} \right\|_{L_x^k L_T^2} \\
&\lesssim \left[T^{\frac{1}{2}} \|u\|_{X_T}^{k} + (1+T^{\frac{1}{4}} \|u\|_{X_T}) \|u\|_{X_T}^{k-1} \|P_{\gg 1} u\|_{X_T} \right] \|P_N u\|_{L_x^k L_T^{\infty}}.
\end{aligned}
$$

Thus, the first term in (5.29) can be bounded by

$$
\lesssim T^{\frac{1}{2}} \|u\|_{X_T}^{k} \|P_{\gg 1} u\|_{X_T} + (1+T^{\frac{1}{4}} \|u\|_{X_T}) \|u\|_{X_T}^{k-1} \|P_{\gg 1} u\|_{X_T}^2.
$$

By the Hölder inequality, we get

$$
\begin{aligned}
\|B_{N,4}\|_{L_x^{\frac{1}{1-\varepsilon}} L_T^2} &\lesssim \|P_{\ll N} u\|_{L_x^{\frac{k(k-2)}{k(1-\varepsilon)-2}} L_T^{\infty}}^{k-2} \|P_N u\|_{L_x^k L_T^{\infty}} \left\| P_{\ll N} u \overline{P_{\ll N} u_x} \right\|_{L_x^k L_T^2} \\
&\lesssim \left[T^{\frac{1}{2}} \|u\|_{X_T}^{k} + (1+T^{\frac{1}{4}} \|u\|_{X_T}) \|u\|_{X_T}^{k-1} \|P_{\gg 1} u\|_{X_T} \right] \|P_N u\|_{L_x^k L_T^{\infty}}.
\end{aligned}
$$

Noticing that $B_{N,4}$ has Fourier support in $|\xi| \sim N$, we can repeat the procedure which we use to deal with the second term in (5.17), and obtain that the second term in (5.29) can be bounded by

$$
\lesssim T^{\frac{1}{2}} \|u\|_{X_T}^{2k} \|P_{\gg 1} u\|_{X_T} + (1+T^{\frac{1}{4}} \|u\|_{X_T}) \|u\|_{X_T}^{2k-1} \|P_{\gg 1} u\|_{X_T}^2.
$$

Therefore, we obtain

$$\begin{aligned}&\left(\sum_{N\gg 1}\left\|\int_0^t S(t-\tau)I_{N,4}(\tau)d\tau\right\|_{Y_T}^2\right)^{1/2}\\ \lesssim&(1+\|u\|_{X_T}^k)\left[T^{\frac{1}{2}}\|u\|_{X_T}^k\|P_{\gg 1}u\|_{X_T}+(1+T^{\frac{1}{4}}\|u\|_{X_T})\|u\|_{X_T}^{k-1}\|P_{\gg 1}u\|_{X_T}^2\right].\end{aligned}$$

5.5. **Nonlinear estimates of $I_{N,5}$.** From (3.5), (3.11), (3.6) and (3.7), we have

$$\begin{aligned}&\left(\sum_{N\gg 1}\left\|\int_0^t S(t-\tau)e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy}B_{N,5}d\tau\right\|_{Y_T}^2\right)^{1/2}\\ \lesssim&\left(\sum_{N\gg 1}\|B_{N,5}\|_{L_x^1L_T^2}^2\right)^{1/2}\\ &+\left(\sum_{N\gg 1}\left(\sum_M\left\|P_M(e^{-\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy}B_{N,5})\right\|_{L_x^1L_T^2}\right)^2\right)^{1/2},\end{aligned}\tag{5.30}$$

where $B_{N,5}=|P_{\ll N}u|^{2k}P_Nu$. By the Hölder inequality, we have

$$\begin{aligned}\|B_{N,5}\|_{L_x^1L_T^2}\lesssim&T^{\frac{1}{2}}\|P_{\ll N}u\|_{L_x^{4k}L_T^\infty}^{2k}\|P_Nu\|_{L_x^2L_T^\infty}\\ \lesssim&T^{\frac{1}{2}}\|u\|_{X_T}^{2k}\|P_Nu\|_{L_x^2L_T^\infty},\end{aligned}$$

and

$$\begin{aligned}\|B_{N,5}\|_{L_x^{\frac{1}{1-\varepsilon}}L_T^2}\lesssim&T^{\frac{1}{2}}\|P_{\ll N}u\|_{L_x^{\frac{4k}{1-2\varepsilon}}L_T^\infty}^{2k}\|P_Nu\|_{L_x^2L_T^\infty}\\ \lesssim&T^{\frac{1}{2}}\|u\|_{X_T}^{2k}\|P_Nu\|_{L_x^2L_T^\infty}.\end{aligned}$$

Thus, in a similar way as dealing with $I_{N,1}$ and $I_{N,4}$, and noticing that $B_{N,5}$ has Fourier support in $|\xi|\sim N$, we can bound (5.30) by

$$\lesssim T^{\frac{1}{2}}\|u\|_{X_T}^{3k}\|P_{\gg 1}u\|_{X_T}.$$

6. A priori estimates for solutions.

By the scaling argument

$$u(t,x)\mapsto u_\gamma(t,x)=\frac{1}{\gamma^{1/k}}u(\frac{t}{\gamma^2},\frac{x}{\gamma}),$$

we have

$$\begin{aligned}\|u_{0,\gamma}\|_{L^2}=&\gamma^{\frac{1}{2}-\frac{1}{k}}\|u_0\|_{L^2},\\ \|u_{0,\gamma}\|_{\dot H^{\frac{1}{2}}}=&\frac{1}{\gamma^{1/k}}\|u_0\|_{\dot H^{\frac{1}{2}}}.\end{aligned}$$

Thus, we may rescale

$$\begin{aligned}\left\|P_{\lesssim 1}u_{0,\gamma}\right\|_{L^2}\leqslant&\gamma^{\frac{1}{2}-\frac{1}{k}}\|u_0\|_{L^2}=C_{low},\\ \|P_{\gg 1}u_{0,\gamma}\|_{H^{\frac{1}{2}}}\leqslant&\frac{1}{\gamma^{1/k}}\|u_0\|_{H^{\frac{1}{2}}}<C_{high}\ll 1,\end{aligned}$$

where we choose $\gamma=\gamma(\|u_0\|_{H^{1/2}})\gg 1$, and take the time interval T depending on γ later. We now drop the writing of the scaling parameter γ and assume

$$\left\|P_{\lesssim 1}u_0\right\|_{L^2}\leqslant C_{low},$$

$$\|P_{\gg 1}u_0\|_{H^{\frac{1}{2}}} \leqslant C_{high} \ll 1.$$

We now apply this to the norms X_T and $H^{1/2}$, and define new version of the norms of X_T and $H^{1/2}$, given by with the decomposition $I = P_{\lesssim 1} + P_{\gg 1}$,

$$\|u\|_{\tilde{X}_T} = \frac{1}{C_{low}}\left\|P_{\lesssim 1}u\right\|_{X_T} + \frac{1}{C_{high}}\|P_{\gg 1}u\|_{X_T},$$

and

$$\|\phi\|_{\tilde{H}^{1/2}} = \frac{1}{C_{low}}\left\|P_{\lesssim 1}\phi\right\|_{L^2} + \frac{1}{C_{high}}\|P_{\gg 1}\phi\|_{H^{1/2}},$$

which implies that $\|u_0\|_{\tilde{H}^{1/2}} \leqslant 2$.

For the low frequency part, we have the following estimates.

Lemma 6.1. *Let u be a solution of* (1.1)-(1.2). *Then*

$$\left\|P_{\lesssim 1}u\right\|_{X_T} \lesssim C_{low} + T^{1/2}\|u\|_{X_T}^{k+1}.$$

Proof. Using the integral equation of (1.1)

$$u(t) = S(t)u_0 - \lambda\int_0^t S(t-\tau)|u(\tau)|^k u_x(\tau)d\tau,$$

and by (3.1), (3.2), (3.3), (3.8), (3.9), (3.10) and the Hölder inequality, we have

$$\begin{aligned}\left\|P_{\lesssim 1}u\right\|_{X_T} &\lesssim \left\|S(t)P_{\lesssim 1}u_0\right\|_{X_T} + \left\|\int_0^t S(t-\tau)P_{\lesssim 1}(|u|^k u_x)(\tau)d\tau\right\|_{X_T} \\ &\lesssim \left\|P_{\lesssim 1}u_0\right\|_{L^2} + \left\|P_{\lesssim 1}(|u|^k u_x)\right\|_{L_T^1 H_x^{1/2}} \lesssim C_{low} + \left\||u|^k u_x\right\|_{L_T^1 L_x^2} \\ &\lesssim C_{low} + T^{1/2}\|u\|_{L_x^{2k}L_T^\infty}^k \|u_x\|_{L_x^\infty L_T^2} \\ &\lesssim C_{low} + T^{1/2}\|u\|_{X_T}^{k+1},\end{aligned}$$

which is the desired result. □

For the high frequency part, we have

Lemma 6.2. *Let u and v_N be given in* (2.1). *Then*

$$\|P_{\gg 1}u\|_{X_T} \lesssim (1 + \|u\|_{L_T^\infty H_x^{1/2}}^{2k})\left(\sum_{N\gg 1}\|v_N\|_{Y_T}^2\right)^{1/2}.$$

Proof. By (2.1), we have

$$P_N u = e^{\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k dy} v_N.$$

For $L_T^\infty H_x^{1/2}$-norm, by the interpolation theorem, we obtain for $N \gg 1$,

$$\begin{aligned}\|P_N u\|_{H_x^{1/2}} &\lesssim \|P_N u\|_{L^2}^{\frac{1}{2}}\|P_N u\|_{H^1}^{\frac{1}{2}} \lesssim \|v_N\|_{L^2}^{\frac{1}{2}}(\|P_N u\|_{L^2} + \|\partial_x P_N u\|_{L^2})^{\frac{1}{2}} \\ &\lesssim \|v_N\|_{L^2}^{\frac{1}{2}}\left(\|v_N\|_{L^2} + \left\||P_{\ll N}u|^k v_N\right\|_{L^2} + \|\partial_x v_N\|_{L^2}\right)^{\frac{1}{2}} \\ &\lesssim \|v_N\|_{L^2}^{\frac{1}{2}}\left(\|P_{\ll N}u\|_{L_x^{4k}}^k\|v_N\|_{L_x^4} + \|v_N\|_{H^1}\right)^{\frac{1}{2}} \\ &\lesssim \left(1 + \|P_{\ll N}u\|_{H_x^{1/2}}^k\right)^{\frac{1}{2}}\|v_N\|_{H_x^{1/2}} \lesssim \left(1 + \|P_{\ll N}u\|_{H_x^{1/2}}^k\right)\|v_N\|_{H_x^{1/2}},\end{aligned}$$

which yields the desired estimate by summing on l_N^2.

For the $L^\infty_x L^2_T$-norm, noticing that

$$\partial_x P_N u = e^{\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k\,dy}\left(\partial_x v_N + \frac{i\lambda}{2}|P_{\ll N}u|^k v_N\right),$$

we have

$$\begin{aligned}\|\partial_x P_N u\|_{L^\infty_x L^2_T} \lesssim{}& \|\partial_x v_N\|_{L^\infty_x L^2_T}\\ &+\left\|\tilde{P}_N\left(\sum_{N_1} P_{N_1}(e^{\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N}u|^k\,dy})|P_{\ll N}u|^k\sum_{N_2} P_{N_2}v_N\right)\right\|_{L^\infty_x L^2_T}.\end{aligned}\tag{6.1}$$

To estimate the second term (6.1), we split the sum $\sum_{N_2}=\sum_{N_2\sim N}+\sum_{N_2\nsim N}$. For $N_2\sim N$, from the Bernstein inequality, we bound (6.1) by

$$\begin{aligned}&\lesssim\left\||P_{\ll N}u|^k\sum_{N_2\sim N}P_{N_2}v_N\right\|_{L^\infty_x L^2_T}\lesssim\|P_{\ll N}u\|^k_{L^\infty_{x,T}}\sum_{N_2\sim N}\|P_{N_2}v_N\|_{L^\infty_x L^2_T}\\ &\lesssim N\left\|D_x^{-1/k}P_{\ll N}u\right\|^k_{L^\infty_{x,T}}\sum_{N_2\sim N}\|P_{N_2}v_N\|_{L^\infty_x L^2_T}\\ &\lesssim\|P_{\ll N}u\|^k_{L^\infty_T H^{1/2}_x}\sum_{N_2\sim N}\|P_{N_2}\partial_x v_N\|_{L^\infty_x L^2_T}\\ &\lesssim\|u\|^k_{L^\infty_T H^{1/2}_x}\sum_{N_2\sim N}\|P_{N_2}\partial_x v_N\|_{L^\infty_x L^2_T}.\end{aligned}$$

For the part $N_2\nsim N$, we split it as $\sum_{N_2\nsim N}=\sum_{N_2\ll N}+\sum_{N_2\gg N}$. Noticing that for $N_2\ll N$,

$$\begin{aligned}&\tilde{P}_N\left(\sum_{N_1}P_{N_1}(e^{\frac{i\lambda}{2}\int_{-\infty}^x|P_{\ll N}u|^k\,dy})|P_{\ll N}u|^k\sum_{N_2}P_{N_2}v_N\right)\\ =&\tilde{P}_N\left(P_{\sim N}(e^{\frac{i\lambda}{2}\int_{-\infty}^x|P_{\ll N}u|^k\,dy})|P_{\ll N}u|^k P_{\ll N}v_N\right),\end{aligned}\tag{6.2}$$

and for $N_2\gg N$,

$$(6.2)=\tilde{P}_N\left(\sum_{N_1\sim N_2\gg N}P_{N_1}(e^{\frac{i\lambda}{2}\int_{-\infty}^x|P_{\ll N}u|^k\,dy})|P_{\ll N}u|^k P_{N_2}v_N\right),$$

we can bound (6.1), in view of the Bernstein inequality and the Hölder inequality, by

$$\begin{aligned}\lesssim&\left\|P_{\sim N}e^{\frac{i\lambda}{2}\int_{-\infty}^x|P_{\ll N}u|^k\,dy}\right\|_{L^\infty_{x,T}}\|P_{\ll N}u\|^k_{L^\infty_{x,T}}\|P_{\ll N}v_N\|_{L^\infty_{x,T}}\\ &+\sum_{N_1\sim N_2\gg N}\left\|P_{N_1}e^{\frac{i\lambda}{2}\int_{-\infty}^x|P_{\ll N}u|^k\,dy}\right\|_{L^\infty_{x,T}}\|P_{\ll N}u\|^k_{L^\infty_{x,T}}\|P_{N_2}v_N\|_{L^\infty_{x,T}}\\ \lesssim&N^{-\frac{k}{(2k+1)}}\left\|P_{\sim N}\partial_x e^{\frac{i\lambda}{2}\int_{-\infty}^x|P_{\ll N}u|^k\,dy}\right\|_{L^\infty_{x,T}}\\ &\quad\cdot\left\|D_x^{-\frac{1}{(2k+1)}}P_{\ll N}u\right\|^k_{L^\infty_{x,T}}\left\|D_x^{-\frac{1}{(2k+1)}}P_{\ll N}v_N\right\|_{L^\infty_{x,T}}\\ &+\sum_{N_1\sim N_2\gg N}N_1^{-\frac{k}{(2k+1)}}\left\|P_{N_1}\partial_x e^{\frac{i\lambda}{2}\int_{-\infty}^x|P_{\ll N}u|^k\,dy}\right\|_{L^\infty_{x,T}}\end{aligned}$$

$$
\begin{aligned}
&\cdot \left\| D_x^{-\frac{1}{(2k+1)}} P_{\ll N} u \right\|^k_{L^\infty_{x,T}} \left\| D_x^{-\frac{1}{(2k+1)}} P_{N_2} v_N \right\|_{L^\infty_{x,T}} \\
\lesssim & N^{-\frac{k}{(2k+1)}} \left\| P_{\ll N} u \right\|^k_{L^\infty_{x,T}} \|u\|^k_{L^\infty_T H^{1/2}_x} \|v_N\|_{L^\infty_T H^{1/2}_x} \\
&+ \sum_{N_1 \sim N_2 \gg N} N_1^{-\frac{k}{3(2k+1)}} N_2^{-\frac{k}{3(2k+1)}} N^{-\frac{k}{3(2k+1)}} \left\| P_{\ll N} u \right\|^k_{L^\infty_{x,T}} \|u\|^k_{L^\infty_T H^{1/2}_x} \|P_{N_2} v_N\|_{L^\infty_T H^{1/2}_x} \\
\lesssim & \|u\|^{2k}_{L^\infty_T H^{1/2}_x} \|v_N\|_{L^\infty_T H^{1/2}_x} .
\end{aligned}
$$

Therefore, summing on l^2_N, we complete the proof for the $L^\infty_x L^2_T$-norm.

For the $L^2_x L^\infty_T$-norm, it is easy to obtain the desired result since $|P_N u| = |v_N|$.

We turn to estimate the $L^4_x L^\infty_T$-norm. It is similar with the proof for the $L^\infty_x L^2_T$-norm, since $\left\| \langle D_x \rangle^{1/4} P_N u \right\|_{L^4_x L^\infty_T} \sim N^{1/4} \|P_N u\|_{L^4_x L^\infty_T}$ for $N \gg 1$. In fact, we have

$$
\left\| \langle D_x \rangle^{1/4} P_N u \right\|_{L^4_x L^\infty_T} \sim N^{1/4} \left\| \sum_{N_1} P_{N_1}\big(e^{\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy}\big) \sum_{N_2} P_{N_2} v_N \right\|_{L^4_x L^\infty_T} . \tag{6.3}
$$

We also split $\sum_{N_2} = \sum_{N_2 \sim N} + \sum_{N_2 \nsim N}$. For $N_2 \sim N$, we bound (6.3) by

$$
\lesssim N^{1/4} \sum_{N_2 \sim N} \|P_{N_2} v_N\|_{L^4_x L^\infty_T} \lesssim \left\| \langle D_x \rangle^{1/4} v_N \right\|_{L^4_x L^\infty_T} .
$$

For the part $N_2 \nsim N$, we split it as $\sum_{N_2 \nsim N} = \sum_{N_2 \ll N} + \sum_{N_2 \gg N}$. Noticing that for $N_2 \ll N$,

$$
\begin{aligned}
&\tilde{P}_N \left(\sum_{N_1} P_{N_1}\big(e^{\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy}\big) \sum_{N_2} P_{N_2} v_N \right) \qquad (6.4) \\
=&\tilde{P}_N \left(P_{\sim N}\big(e^{\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy}\big) P_{\ll N} v_N \right),
\end{aligned}
$$

and for $N_2 \gg N$,

$$
(6.4) = \tilde{P}_N \left(\sum_{N_1 \sim N_2 \gg N} P_{N_1}\big(e^{\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy}\big) P_{N_2} v_N \right),
$$

we can bound (6.3), in view of the Bernstein inequality and the Hölder inequality, by

$$
\begin{aligned}
\lesssim & \left\| \langle D_x \rangle^{1/4} v_N \right\|_{L^4_x L^\infty_T} + N^{1/4} \left\| P_{\sim N}\big(e^{\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy}\big) \right\|_{L^\infty_{x,T}} \|P_{\ll N} v_N\|_{L^4_x L^\infty_T} \\
&+ N^{1/4} \sum_{N_1 \sim N_2 \gg N} \left\| P_{N_1}\big(e^{\frac{i\lambda}{2}\int_{-\infty}^x |P_{\ll N} u|^k dy}\big) \right\|_{L^\infty_{x,T}} \|P_{N_2} v_N\|_{L^4_x L^\infty_T} \\
\lesssim & \left\| \langle D_x \rangle^{1/4} v_N \right\|_{L^4_x L^\infty_T} + N^{-3/4} \|P_{\ll N} u\|^k_{L^\infty_{x,T}} \left\| \langle D_x \rangle^{1/4} P_{\ll N} v_N \right\|_{L^4_x L^\infty_T} \\
&+ \sum_{N_1 \sim N_2 \gg N} N_1^{-1/3} N_2^{-1/3} N^{-1/3} \|P_{\ll N} u\|^k_{L^\infty_{x,T}} \left\| \langle D_x \rangle^{1/4} P_{N_2} v_N \right\|_{L^4_x L^\infty_T} \\
\lesssim & (1 + \|u\|^k_{L^\infty_T H^{1/2}_x}) \left\| \langle D_x \rangle^{1/4} v_N \right\|_{L^4_x L^\infty_T} ,
\end{aligned}
$$

which yields the desired estimate by applying l^2_N-sum.

Thus, we complete the proof of this Lemma. □

Of course, we need the following estimate of the data.

Lemma 6.3. *For any $u_0 \in H^{1/2}$, we have*

$$\left(\sum_{N\gg 1}\left\|S(t)(e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u_0|^k\,dy}P_N u_0)\right\|_{Y_T}^2\right)^{1/2} \lesssim (1+\|u_0\|_{H^{1/2}}^k)\,\|P_{\gg 1}u_0\|_{H^{1/2}}\,. \tag{6.5}$$

Proof. From (3.1), (3.2) and (3.3), we bound the left hand side of (6.5) by

$$\lesssim \left(\sum_{N\gg 1}\left\|e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u_0|^k\,dy}P_N u_0\right\|_{H^{1/2}}^2\right)^{1/2} \tag{6.6}$$

$$+\left(\sum_{N\gg 1}\left(\sum_{M}\left\|P_M(e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u_0|^k\,dy}P_N u_0)\right\|_{H^{1/2}}\right)^2\right)^{1/2}. \tag{6.7}$$

From Lemma 3.3, we have

$$\begin{aligned}(6.6) \lesssim{}& \left(\sum_{N\gg 1}[\|P_{\ll N}u_0\|_{L^{4k}}^{2k}\,\|P_N u_0\|_{L^4}^2 + \|P_N u_0\|_{H^{1/2}}^2]\right)^{1/2}\\ \lesssim{}&(1+\|u_0\|_{H^{1/2}}^2)\,\|P_{\gg 1}u_0\|_{H^{1/2}}\,.\end{aligned}$$

For the second term (6.7), it is similar with (5.21). We split the sum $\sum_M = \sum_{M\lesssim N} + \sum_{M\gg N}$. By the Bernstein inequality, the Hölder inequality and the Sobolev embedding theorem, we bound (6.7) by

$$\begin{aligned}
\lesssim{}& \left(\sum_{N\gg 1}\left(\sum_{M\lesssim N}\langle M\rangle^{\frac12}\,\|P_N u_0\|_{L_x^2}\right)^2\right)^{1/2}\\
&+\left(\sum_{N\gg 1}\left(\sum_{M\gg N}\left\|\langle D_x\rangle^{\frac12}P_M(e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u_0|^k\,dy}P_N u_0)\right\|_{L_x^2}\right)^2\right)^{1/2}\\
\lesssim{}& \left(\sum_{N\gg 1}\left(N^{1/2}\,\|P_N u_0\|_{L_x^2}\right)^2\right)^{1/2}\\
&+\left(\sum_{N\gg 1}\left(\sum_{M\gg N}M^{\frac12}\left\|(P_{\sim M}e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u_0|^k\,dy})P_N u_0\right\|_{L_x^2}\right)^2\right)^{1/2}\\
\lesssim{}& \|P_{\gg 1}u_0\|_{H^{1/2}}\\
&+\left(\sum_{N\gg 1}\left(\sum_{M\gg N}M^{\frac12}\left\|P_{\sim M}e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u_0|^k\,dy}\right\|_{L^4}\|P_N u_0\|_{L_x^4}\right)^2\right)^{1/2}\\
\lesssim{}& \|P_{\gg 1}u_0\|_{H^{1/2}}\\
&+\left(\sum_{N\gg 1}\left(\sum_{M\gg N}M^{-\frac12}\left\|P_{\sim M}\partial_x e^{-\frac{i\lambda}{2}\int_{-\infty}^{x}|P_{\ll N}u_0|^k\,dy}\right\|_{L^4}\|P_N u_0\|_{L_x^4}\right)^2\right)^{1/2}
\end{aligned}$$

$$\begin{aligned}
&\lesssim \|P_{\gg 1}u_0\|_{H^{1/2}} + \left(\sum_{N\gg 1}\left(\|P_{\ll N}u_0\|_{L^{4k}}^{k}\|P_N u_0\|_{L_x^4}\right)^2\right)^{1/2}\\
&\lesssim (1+\|u_0\|_{H^{1/2}}^{k})\|P_{\gg 1}u_0\|_{H^{1/2}},
\end{aligned}$$

which yields the desired result. □

With the help of the above lemmas, we can prove the following proposition which yields the a priori estimate.

Proposition 6.1. *Let u be a smooth solution to* (1.1)-(1.2) *and* $0 < T \leqslant C_{high}^4$. *Then we have*

$$\|u\|_{\tilde{X}_T} \leqslant C(C_{low}) + C(C_{low} + \|u\|_{\tilde{X}_T})^{3k}(T^{1/4} + C_{high})\|u\|_{\tilde{X}_T}.$$

Proof. Noticing that

$$\left\|P_{\lesssim 1}u\right\|_{X_T} \lesssim C_{low}\|u\|_{\tilde{X}_T}, \quad \|P_{\gg 1}u\|_{X_T} \lesssim C_{high}\|u\|_{\tilde{X}_T}.$$

and from Lemmas 6.1, 6.2, 6.3 and Proposition 5.1, we obtain through a complicated computation

$$\begin{aligned}
&\|u\|_{\tilde{X}_T} = \frac{1}{C_{low}}\left\|P_{\lesssim 1}u\right\|_{X_T} + \frac{1}{C_{high}}\|P_{\gg 1}u\|_{X_T}\\
\lesssim& 1 + \frac{1}{C_{low}}T^{\frac12}\|u\|_{X_T}^{k+1} + \frac{1}{C_{high}}(1+\|u\|_{L_T^\infty H_x^{1/2}}^{2k})\left(\sum_{N\gg 1}\|v_N\|_{Y_T}^2\right)^{1/2}\\
\lesssim& 1 + \frac{1}{C_{low}}T^{\frac12}\|u\|_{X_T}^{k+1} + \frac{1}{C_{high}}(1+\|u\|_{L_T^\infty H_x^{1/2}}^{2k})(1+\|u_0\|_{H^{1/2}}^{k})\|P_{\gg 1}u_0\|_{H^{1/2}}\\
&+\frac{1}{C_{high}}(1+\|u\|_{L_T^\infty H_x^{1/2}}^{2k})\Big\{T^{\frac12}\|u\|_{X_T}^{3k}\|P_{\gg 1}u\|_{X_T}\\
&+(1+\|u\|_{X_T}^{k})\left[T^{\frac12}\|u\|_{X_T}^{\tilde{k}+1}\|P_{\gg 1}u\|_{X_T}^{k-\tilde{k}} + (1+T^{\frac14}\|u\|_{X_T})\|u\|_{X_T}^{\tilde{k}}\|P_{\gg 1}u\|_{X_T}^{k+1-\tilde{k}}\right]\\
&+T^{\frac12}\left(\|u\|_{X_T}^{2k-1} + \|u\|_{X_T}^{(5k-2)/2}\right)\|P_{\gg 1}u\|_{X_T}\\
&+T^{\frac14}(1+T^{\frac14}\|u\|_{X_T})^{\frac12}\|u\|_{X_T}^{2k-1}\|P_{\gg 1}u\|_{X_T}^{\frac32}\\
&+(1+\|u\|_{X_T}^{k})\left[T^{\frac12}\|u\|_{X_T}^{k}\|P_{\gg 1}u\|_{X_T} + (1+T^{\frac14}\|u\|_{X_T})^2\|u\|_{X_T}^{k-1}\|P_{\gg 1}u\|_{X_T}^2\right]\Big\}\\
\leqslant& C + C(C_{low})T^{\frac12}\|u\|_{\tilde{X}_T}^{k+1} + (1+\|u\|_{L_T^\infty H_x^{1/2}}^{2k})(1+\|u_0\|_{H^{1/2}}^{k})\\
&+C(C_{low})(1+\|u\|_{L_T^\infty H_x^{1/2}}^{2k})\Big\{T^{\frac14}\|u\|_{\tilde{X}_T}^{3k+1} + T^{\frac14}(1+\|u\|_{\tilde{X}_T}^{k})\|u\|_{\tilde{X}_T}^{k+1}\\
&+T^{\frac12}(\|u\|_{\tilde{X}_T}^{k+1} + \|u\|_{\tilde{X}_T}^{5k/2}) + C_{high}^{k-\tilde{k}}(1+T^{\frac14}\|u\|_{\tilde{X}_T})(1+\|u\|_{\tilde{X}_T}^{k})\|u\|_{\tilde{X}_T}^{k+1}\\
&+C_{high}^{3/2}(1+T^{\frac14}\|u\|_{\tilde{X}_T})\|u\|_{\tilde{X}_T}^{2k} + C_{high}^{2}(1+T^{\frac14}\|u\|_{\tilde{X}_T})^2(1+\|u\|_{\tilde{X}_T}^{k})\|u\|_{\tilde{X}_T}^{k+1}\Big\}.
\end{aligned}$$

Notice that

$$\|u(t)\|_{H^{1/2}} \lesssim \left\|P_{\lesssim 1}u(t)\right\|_{L^2} + C_{high}\|P_{\gg 1}\|_{\tilde{H}^{1/2}}.$$

The high frequency part $C_{high}\|P_{\gg 1}\|_{\tilde{H}^{1/2}}$ can be absorbed into the $\tilde{X}_T$-norm. Then substituting Lemma 6.1 again in estimating the low frequency part of the norm $\left\|P_{\lesssim 1}u\right\|_{L_T^\infty H_x^{1/2}}$, we complete the proof of Proposition 6.1. □

From Proposition 6.1, we have the following a priori estimate for the solution of (1.1)-(1.2) if we take T and C_{high} small enough.

Corollary 6.1. *Let u be a smooth solution to* (1.1)-(1.2). *we have*

$$\|u\|_{\tilde{X}_T} \lesssim C_{low} + C_{high},$$

for T and C_{high} small enough.

For the proof of Theorem 1.1, we can follow the compactness argument with the a priori estimate. Since the proof is standard, we omit the details and refer to the papers [14, 15, 16, 17, 19, 22].

Acknowledgements. The author would like to thank the referees for valuable comments and suggestions on the original manuscript and Prof. L. Hsiao for her frequent encouragement.

REFERENCES

[1] J. Bergh, J. Löfström, Interpolation Spaces, An Introduction, Springer-Verlag, Berlin Heidelberg (1976).

[2] J. Bourgain, *Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations, Parts I, II*, Geom. Funct. Anal. **3** (1993), 107–156, 209–262.

[3] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, T. Tao, *Global well-posedness for the Schrödinger equations with derivative*, SIAM J. Math. Anal. **33** (2001), 649–669.

[4] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, T. Tao, *A refined global well-posedness for the Schrödinger equations with derivative*, SIAM J. Math. Anal. **34** (2002), 64–86.

[5] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, T. Tao, *Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in $\mathbb{R}^3$*, to appear in Annals Math., `arXiv:math.AP/0402129`.

[6] I. Fukuda, Y. Tsutsumi, *On solutions of the derivative nonlinear Schrödinger equation: existence and uniqueness theorem*, Funkcial. Ekvac. **23** (1980), 259–277.

[7] I. Fukuda, Y. Tsutsumi, *On solutions of the derivative nonlinear Schrödinger equation II*, Funkcial. Ekvac. **234** (1981), 85–94.

[8] A. Grüenrock, *On the Cauchy and periodic boundary value problem for a certain class of derivative nonlinear Schrödinger equations*, preprint.

[9] N. Hayashi, *The initial value problem for the derivative nonlinear Schrödinger equation in the energy space*, Nonlinear Anal. **20** (1993), 823–833.

[10] N. Hayashi, T. Ozawa, *On the derivative nonlinear Schrödinger equation*, Phys D **55**(1992), 14–36.

[11] N. Hayashi, T. Ozawa, *Remarks on nonlinear Schrödinger equations in one space dimension*, Diff. Integ. Eq. **7** (1994), 453–461.

[12] C. Kenig, G. Ponce, L. Vega, *Oscillatory integrals and regularity of dispersive equations*, Indiana Univ. Math. J. **40** (1991), 33–69.

[13] C. E. Kenig, G. Ponce, L. Vega, *Small solutions to nonlinear Schrödinger equations*, Ann. Inst. H. Poincaré Anal. Non Lineaire, **10** (1993), 255–288.

[14] C. E. Kenig, H. Takaoka, *Global wellposedness of the modified Benjamin-Ono equation with initial data in $H^{1/2}$*, Int. Math. Res. Not., (2006), Art. ID 95702, 44 pages.

[15] H. Koch, N. Tzvetkov, *On the local well-posedness of the Benjamin-Ono equation in $H^s(\mathbb{R})$*, Int. Math. Res. Not., **26** (2003), 1449–1464.

[16] L. Molinet, F. Ribaud, *Well-posedness results for the generalized Benjamin-Ono equation with small initial data*, J. Math. Pures Appl., **83** (2004), 277–311.

[17] L. Molinet, F. Ribaud, *Well-posedness results for the generalized Benjamin-Ono equation with arbitrary large initial data*, Int. Math. Res. Not., **70** (2004), 3757–3795.

[18] T. Ozawa, *On the nonlinear Schrödinger equations of derivative type*, Indiana Univ. Math. J. **45** (1996), 137–163.

[19] G. Ponce, *On the global well-posedness of the Benjamin-Ono equation*, Diff. Int. Eqns., **4** (1991), 527–542.

[20] E. M. Stein, Harmonic analysis, Real-variable methods, orthogonality, and oscillatory integrals, Princeton University Press, Princeton (1993).

[21] H. Takaoka, *Well-posedness for the one-dimensional nonlinear Schrödinger equation with the derivative nonlinearity*, Adv. Diff. Eqns., **4** (1999), 561-580.

[22] T. Tao, *Global well-posedness of the Benjamin-Ono equation in* $H^1(\mathbb{R})$, J. Hyperbolic Diff. Eqns., **1** (2004), 27–49.

[23] T. Tao, Nonlinear Dispersive Equations: Local and Global Analysis, CBMS Regional Conference Series in Mathematics, **106**, American Mathematical Society, Providence, RI, (2006).

Received July 25, 2006; revised November 2006.

E-mail address: `hcc@amss.ac.cn`